

The critical care nursing role in low and lower-middle income settings: A scoping review

Andy Macey^{1,2}, Gerard O'Reilly^{1,3,4}, Ged Williams⁵, Peter Cameron¹

Corresponding author: Andy Macey

6 Ostend St. Bittern, VIC, 3918

Andrew.macey@monash.edu

+61 0413 063 233

Mr. Andy Macey *CCRN M.Ed.*

¹ Monash University, School of Public Health and Preventative Medicine, Melbourne, Australia

² Peninsula Health, The Learning Hub, Melbourne, Australia

A/Prof. Gerard O'Reilly *MBBS MPH MBiostat AStat FACEM PhD*

¹ Monash University, School of Public Health and Preventative Medicine, Melbourne, Australia

³ The Alfred, Emergency and Trauma Centre, Melbourne, Australia

⁴ The Alfred, National Trauma Research Institute, Melbourne, Australia

Prof. Ged Williams *RN, FACN, FACHSM, FAAN*

⁵ Griffith University, School of Nursing and Midwifery, Brisbane, Australia

Prof. Peter Cameron *MBBS, MD, FACEM, FRCEM (hon)*

¹ Monash University, School of Public Health and Preventative Medicine, Melbourne, Australia

Key words:

- Critical Care
- Low- and Middle-Income Countries
- Nursing
- Delivery of Healthcare
- Health Care Facilities, Manpower, and Services

Abstract

Objectives

A scoping review was conducted to answer the question: How is critical care nursing (CCN) performed in low-income and lower-middle income countries (LICs/LMICs)?

Design

Scoping review guided by the JBI Manual for Evidence Synthesis

Data sources

Six electronic databases and five web-based resources were systematically searched to identify relevant literature published between 2010 - April 2021.

Review methods

The search results received two stage screening: 1) Title and abstract 2) Full text screening. For sources of evidence to progress, agreement needed to be reached by two reviewers. Data were extracted and cross checked. Data were analysed, sorted by themes, and mapped to region and country.

Results

Literature was reported across five geo-regions. Nurses with a range formal and informal training were identified as providing critical care. Availability of staff was frequently reported as a problem. No reports provided a comprehensive description of CCN in LICs/LMICs. However, a variety of nursing practices and non-clinical responsibilities were highlighted. Availability of equipment to fulfil the nursing role were widely discussed. Perceptions of inadequate resourcing were common. Under- and post-graduate level preparation were poorly described but frequently reported. The delivery of short format

critical care courses was more fully described. There were reports of educational evaluation, especially regarding internationally supported initiatives.

Conclusions

Despite commonalities, CCN is unique to regional and socio-economic contexts. Nurses work within a complex team, yet the structure and skill levels of such teams will vary according to patient population, resources, and treatments available. Therefore, a universal definition of the CCN role in LIC/LMIC health systems is likely unhelpful. Research to elucidate current assets, capacity and needs of nurses providing critical care in specific LIC/LMIC contexts is needed. Outputs from such research would be invaluable in supporting contextually appropriate capacity development programs.

Strengths and limitations of this review

- This is the first review to explore and map the evidence related to how CCN is performed in LICs/LMICs
- Literature from all geo-regions where LICs/LMICs are situated was considered for inclusion.
- The inability to translate non-English language documents, mean it is possible that some literature was not captured.
- A lack of literature from Latin American LICs/LMICs in English prevents an accurate picture of critical care nursing in these contexts to be drawn.

Introduction

The development of critical care services in low-income and lower-middle-income countries (LICs/LMICs) is increasing.[1] However, little is known regarding capacity and resourcing of these services.[2] Nurses are key in the successful delivery of critical care in any health-system,[3] yet it is unclear what is currently known regarding their role in LIC/LMIC health systems.

Background

The burden of critical illness is substantial in LICs/LMICs[2, 4, 5] and outcomes are often reported as poorer than in high-resource contexts.[5] The availability of resources to manage critically ill patients in LICs/LMICs are commonly restricted when compared to high-income countries (HICs).[6] Access to high-cost technologies and highly trained, specialised teams of healthcare staff available in high-resource health-systems[1] are not always available in a resource-limited setting.[7] Therefore, the organisation of critical care services needs to be context-specific[8] and will vary across socio-economic and geo-regional situations.

Nurses are the largest occupational group of healthcare workers globally and are central to meeting the sustainable development goals (SDGs) and universal health coverage (UHC).[9] Positive associations between the numbers of nurses employed to provide patient care, their level of education, and improved patient outcomes is well documented in several HICs,[10] and may be amplified in critical care settings.[11] In response, standards recognising the need for nursing models specific to critical care, extended organisational supports, and specialised training to fulfil the critical care nursing (CCN) role are common in HIC's.[11, 12]

Furthermore, the roles and responsibilities of critical care nurses are well described in many HICs. For example, in Australia, nurses routinely undertake all technical and non-technical care of the critically ill patient. They operate and manage therapies such as mechanical

ventilation, extracorporeal therapy, and intra-aortic balloon pumps. Additionally, they might measure cardiac output from highly technical hemodynamic devices and titrate vasoactive drugs in a semi-autonomous fashion.[13] To date, little work comprehensively reporting the roles, practices, and training of critical care nurses in LICs/LMICs appears to be available. Therefore, a scoping review was conducted to map the available evidence, identify the main concepts, sources of data and knowledge gaps and answer the question: How is the CCN role performed in LICs/LMICs?

Objectives

To elucidate the role of nurses working in critical care contexts in LICs/LMICs the review aimed to address four sub-questions:

- 1) Who is involved in providing CCN in LICs/LMICs?
- 2) What are the day-to-day roles and responsibilities of nurses providing critical care in LICs/LMICs?
- 3) What professional scope of practice is reported for nurses providing critical care in LICs/LMICs?
- 4) What training and ongoing education are available to nurses providing critical care in LICs/LMICs?

Methods

The scoping review was guided by the JBI Manual for Evidence Synthesis[14] and the Preferred Reporting Items for Systematic Reviews extension for scoping reviews.[15] A protocol was developed *a priori*, published prospectively, and can be viewed on the Monash University repository for research data.[16]

Search methods

A Population Concept Context (PCC) framework was used to define the evidence to be included in this review (Table 1).

Table 1: Population Concepts Context (PCC) framework used to define the evidence included in this review

	<i>Inclusions</i>	<i>Exclusions</i>
Population	Nurses providing critical care or their proxies*.	Visiting international nurses engaged in capacity development programs
Concepts	CCN roles, responsibilities, scope of practices, training, and education.	
Contexts	Low income and lower middle income countries (LICs/LMICs) as defined by the World Bank Country and Lending Groups.[1]	Military hospitals The pre-hospital setting or during the intra-hospital transport of patients

* Proxies for nurses providing critical care include health care workers and lay persons who are not nurses but provide care that might be associated with the nursing role. Examples might include respiratory therapists or healthcare assistants.

1. The World Bank Group. World Bank Country and Lending Groups 2020 [Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>].

Preliminary searches of Ovid Medline, Embase, and Emcare were performed to test and refine the search strategy. Final searches were completed on the 30th of July 2020 and updated on the 7th of April 2021. Six databases were included: Medline, Embase, Emcare, Global Health, SCOPUS, and Web of Science. Limits set in all database searches were publications in the last 10 years (2010-current) and publications in English language. Full details of the search strings employed, additional limits in individual databases, and database versions can be found in the supplementary file 1. Results from all searches were exported to Endnote™ citation management software and then imported into Covidence™ review management software.

Web based searches were completed on 30th July 2020. Five resources were accessed: Google Scholar™, Google™, The World Health Organisation (WHO) website, The World Federation of Critical Care Nurses (WFCCN) website, and websites of CCN organisations listed as members of the WFCCN. The search in Google scholar™ was limited to results published in the last ten years. Consistent with the approach suggested by Pham et al[17] the first 100 search results were submitted to an initial two stage screening process: All titles considered relevant to the research questions were exported to Endnote™. Individual titles were resubmitted to Google Scholar™, accessed, and categorised by type. Evidence that was unrelated to the research questions was removed at this stage. The remaining sources of evidence were imported into Covidence™ for title and abstract screening.

Keyword searching within the WHO website was undertaken using Google™. The WFCCN website was accessed, and an additional 11 regional critical care nursing societies websites were identified for exploration and data extraction as tier-3 sources. The first 100 search results in google were screened for apparent relevance to the research questions. Duplicate website links and results already discovered through searches were ignored.

Search outcome

A total of 361 papers were retrieved from database and web-based searching. Two additional papers were added by hand searching the reference lists of included papers. 31 duplicates were removed. 17 web-based sources of evidence were identified.

Quality appraisal

Guided by Kepes' taxonomy of sources of samples,[18] two tiers of evidence were considered for inclusion in the screening process: Tier-1, the peer reviewed literature. Tier-2, grey literature produced by a range of bodies such as governments or non-commercial organisations, including conference papers, dissertations, and reports. Additionally, Tier-3


sources were considered for separate analysis. Tier-3 samples included commercial and personal websites, blogs, and social media. As is conventional in such a broad scoping review, no critical appraisal of the included literature was undertaken.[14]

Data abstraction

The search results received two stage screening: 1) Title and abstract 2) Full text screening.

For sources of evidence to progress, agreement needed to be reached by two reviewers. A PRISMA flow diagram demonstrating the screening and selection of literature for inclusion is presented in figure 1.

Figure 1: PRISMA Flow chart



PRISMA Flow chart detailing the results from searching databases and other sources, the number of abstracts screened, the number of full texts accessed, and the number of papers included in the final synthesis

Synthesis

Initial extraction was undertaken using a data-charting instrument mapped to the research sub-questions. An iterative approach was taken. As new items or themes that provided useful data became apparent this was discussed by the team and added to the instrument if

appropriate. Finally, three extractions were cross-checked by another member of the team to determine agreement that all relevant data had been charted.

Patient and public involvement

No patient involved.

Ethics approval

No ethics approval was required for this research


Results

Data were predominantly found within peer reviewed journal articles (85%). Other sources of data included conference papers (9%), books (3%), thesis (2%), and reports (1%). Half of the included literature reported quantitative research data. Descriptive or descriptive/correlational methods were the most reported methodologies. 16% of included papers reported qualitative results and a further 6% reported the results of mixed-methods research. The remaining literature encompassed reviews (13%), or expert opinion/editorial papers (15%). One book was identified which defied categorisation as it included aspects of narrative review, expert opinion, and case studies.

The literature focussed on countries within five of the seven geo-regions defined by the World Bank.[19] The literature distributed across the five geo-regions were: Sub-Saharan Africa (SSA) (45%), South Asia (SA) (24%), Middle East and North Africa (MENA) (13%), East Asia and the Pacific (EAP) (10%) and Latin America and the Caribbean (LAC) (2%). The remaining 6% of the literature spoke more broadly to CCN in resource limited settings, not specifically within a region.

Several clinical settings were reported as sites where CCN was practiced. The distribution is visualised in Figure 2.

Figure 2: Distribution of locations of CCN practice in LICs/LMICs



The chart shows the distribution of clinical locations of CCN practice in LICs/LMICs found in the included literature

A large amount of the evidence discussed more than one unit within a health service, for example multiple Intensive Care Units (ICUs) or a combination of ICUs, Paediatric ICUs, Neonatal ICUs, and non-traditional critical care settings. The remaining literature provided evidence of practice within individual dedicated adult, paediatric, neonatal, or mixed units where several patient categories were admitted. A small amount of the literature discussed CCN in Emergency Departments. Finally, some papers spoke only to critical care undertaken in non-specialist environments such as wards or community health centres.

Tier-3 sources of data included three broad categories of website: 1) CCN society websites or social media pages ($n=11$), 2) Blogs ($n=3$), and 3) miscellaneous webpages ($n=3$). The websites of CCN societies and federations included local in-country ($n=8$), regional ($n=2$), and global ($n=1$) level groups.

Three regional CCN groups used Facebook™ groups as their online medium. The remaining critical care societies and federations utilised hosted websites. The blog entries proved to all be by one author, a CCN educator providing trainings in Nepal. A link was discovered in one

of the blog entries to an NGO providing respiratory care education in LIC's/LMICs. All three miscellaneous websites provided evidence of a not-for-profit education company providing free CCN online educational content to LIC's/LMICs.

Literature providing answers to the research questions is displayed in tables 2-5. The corresponding references can be found in supplementary file 2. Both positive and negative results were extracted meaning that discussion of a concept being present or not being present were both captured.

Table 2 identifies sources of data related to who provides nursing care to critically ill patients in LICs/LMICs and maps them to region and countries.


Table 2: Literature providing evidence of who is involved in providing nursing care of critical care patients in low income and lower middle-income countries grouped thematically and mapped to region and country

Data	Region	Citations by country
<i>Qualified nurses with basic training or level of qualification identified</i>	SSA	Ethiopia[1], Kenya[2, 3], Malawi[4, 5], Rwanda[6], Tanzania[7], Uganda[8], Zambia[9]
	SA	India[10-12], Nepal[13]
	EAP	Philippines[14-16], Solomon Islands[17], Vietnam[18]
	MENA	Egypt[19-28], Palestine[29]
	LAC	None
<i>Qualified nurses with post-basic training or level of qualification identified</i>	SSA	Ghana[30], Kenya[2, 3, 31-33], Malawi[4], Mozambique[34], Rwanda[6, 35], Tanzania[7, 36, 37], Uganda[38], Zambia[39], Zimbabwe[40]
	SA	India[10-12, 41], Sri Lanka[42-44]
	EAP	Cambodia[45], Philippines[14, 16]
	MENA	None
	LAC	Haiti [46]
<i>Other nursing cadres*</i>	SSA	Kenya[47], Malawi[4, 5, 48], Mozambique[40, 49], Zimbabwe[40, 49]
	SA	None
	EAP	Philippines[14, 15, 50], Solomon Islands[17]
	MENA	Egypt[28, 51]
	LAC	None
<i>Non-nursing healthcare workers[#]</i>	SSA	Kenya[32, 52], Tanzania[7, 36, 53, 54], Uganda[38]
	SA	India[41, 55], Pakistan[56], Sri Lanka[42]
	EAP	Philippines[16]

	MENA	Egypt[57]
	LAC	None
Non-healthcare trained persons[§]	SSA	Malawi[48], Rwanda[58], Tanzania[59], Zambia[39]
	SA	India[10]
	EAP	None
	MENA	None
	LAC	None
<p>* Other nursing cadres may include but are not limited to student nurses or midwives. # Non-nursing healthcare workers may include but are not limited to clinical officers, respiratory therapists, or ancillary staff. [§] Non-healthcare trained persons may include but are not limited to families and laypersons SSA: Sub Saharan Africa, SA: South Asia, EAP: East Asia and the Pacific, Mena: Middle East and North Africa, LAC: Latin America and the Caribbean The corresponding references for this table are included in supplementary file 2</p>		

Most documents indicated that nurses identified as registered or qualified were engaged in the care of critically ill patients within each region. Many also referred to basic or post-basic qualifications. In several cases these were limited to identifying basic or post-basic trained nurses as being present. Whereas others provided the title of the qualifications (Fig 3).


Figure 3: Distribution of the literature identifying the level of qualification of nurses providing critical care in LICs/LMICs



The chart shows the distribution of included papers that report on or discuss the levels of qualification of nurses providing critical care in LICs/LMICs

A variety of other cadres of healthcare workers and lay persons were also identified as providing aspects or the totality of what might be considered CCN for example midwives and respiratory therapists (Fig 4).

Figure 4: Distribution of the literature identifying student nurses, other cadres of healthcare workers, and lay-persons providing nursing care to critically ill patients in LICs/LMICs



The chart shows the distribution of included papers that report on or discuss student nurses, other cadres of healthcare workers, and lay-persons providing nursing care to critically ill patients in LICs/LMICs

Table 3 identifies sources of data related to the day-to-day roles and responsibilities of nurses caring for critically ill patients in LICs/LMICs and maps them to regions and countries.


Table 3: Literature providing evidence of the day-to-day roles and responsibilities providing nursing care of critical care patients in low income and lower middle-income countries grouped thematically and mapped to region and country

Data	Region	Citations by country
The critical care nursing role, practices, or tasks	SSA	Ethiopia [1], Kenya [2], Malawi [3, 4], Mozambique [5], Nigeria [6, 7], Rwanda [8], Tanzania [9-12], Uganda [13-16], Zambia [17-19]
	SA	India[20], Nepal[21, 22], Pakistan[23, 24], Sri Lanka[25]

	EAP	Cambodia[26, 27], Myanmar[28], , Philippines[29, 30], Vietnam[30, 31], Unnamed EAP country[32]
	MENA	Egypt[33-39], Palestine[40]
	LAC	None
<i>Critical care nursing systems/staffing/capacity</i>	SSA	Kenya [2, 41-43], Malawi[3, 44], Rwanda[45], Tanzania[46], Uganda[14, 46], Zambia[17, 18, 47], Zimbabwe[48]
	SA	India[49-51], Nepal[52, 53], Pakistan[24, 54] Sri Lanka[55-57]
	EAP	Myanmar[28], Philippines[29, 30] Solomon Islands[58], Vietnam[29]
	MENA	Egypt[34-36, 39, 59, 60], Palestine[40]
	LAC	None
<i>The availability and utilisation of physical resources or technologies to fulfil the critical care nursing role</i>	SSA	Kenya[2, 41-43, 61], Malawi[3, 44], Nigeria[6], Rwanda[8, 45], Tanzania [9, 62, 63], Uganda[14, 46], Zambia[18, 47], Zimbabwe[48]
	SA	India[50, 51, 64, 65], Nepal[52, 53], Pakistan[24], Sri Lanka[54, 55]
	EAP	Philippines[30], Solomon Islands[58]
	MENA	Egypt[34, 36, 39], Multiple MENA countries[60]
	LAC	None
<i>The availability of policies or guidelines to support evidence-based practice</i>	SSA	Ethiopia[1], Kenya[42], Malawi[3], Nigeria[6, 7], Rwanda[45] Tanzania[9, 10, 63], Uganda[14-16], Zambia[17-19, 47], Zimbabwe[48]
	SA	India[64, 66-68], Nepal[21, 52], Pakistan[24], Sri Lanka[25]
	EAP	Philippines[29], Vietnam[31]
	MENA	Egypt[34, 35, 39], Multiple MENA countries[60]
	LAC	None
<i>SSA: Sub Saharan Africa, SA: South Asia, EAP: East Asia and the Pacific, Mena: Middle East and North Africa, LAC: Latin America and the Caribbean</i>		
<i>The corresponding references for this table are included in supplementary file 2</i>		

Several specific activities and roles performed by nurses providing critical care in LICs/LMICs were able to be extracted. Some examples included management of airways and mechanical ventilation, care of the septic patient and physiological monitoring. A full overview of the distribution of activities is presented in figure 5.

Figure 5: Distribution of the literature identifying CCN practices or tasks in LICs/LMICs



The chart shows the distribution of included papers that report on or discuss CCN practices or tasks in LICs/LMICs

Details relating to CCN systems, staffing, and capacity were able to be extracted from the literature. Thematically these grouped into four categories: 1) identification of nurse-patient ratios $n=17$, 2) national critical care workforce $n=7$, 3) structure, and staffing of unit-based workforce $n=18$, 4) perceptions of insufficient nursing workforce or poor working conditions $n=5$.

Data regarding the availability and utilisation of physical resources to fulfil the CCN role were also extracted and mapped to nine themes: 1) comprehensive lists of available or recommended equipment $n=10$, 2) mechanical ventilators $n=6$, 3) monitoring technologies $n=2$, 4) oxygen or electricity supply $n=1$, 5) neonatal equipment $n=2$, 6) personal protective equipment $n=1$, 7) other therapeutic devices $n=1$, 8) documentation of care $n=5$, 9) a perceived inadequacy of physical resources $n=9$.

Finally, papers discussing the availability of policies or guidelines were extracted and mapped to themes of locally developed practice guidelines $n=14$, national standards, or recommendations for policy $n=4$, the utilisation of international practice guidelines $n=4$, or perceived unavailability or inadequacy of guidelines $n=11$.

Little data was able to be extracted to answer the question: “What professional scope of practice is reported for nurses providing critical care in LICs/LMICs?” (Table 4).

Table 4: Literature providing evidence information on registration and professional scope of nurses providing nursing care of critical care patients in low income and lower middle-income countries grouped thematically and mapped to region and country

Data	Region	Citations by country
Registration or professional standards	SSA	Malawi[1, 2], Mozambique [3], Rwanda[4, 5], Tanzania[6], Uganda[7, 8], Zambia[9], Multiple SSA countries[10]
	SA	Sri Lanka[11]
	EAP	Philippines[12]
	MENA	Egypt[13, 14], Palestine[15]
	LAC	None

SSA: Sub Saharan Africa, SA: South Asia, EAP: East Asia and the Pacific, Mena: Middle East and North Africa, LAC: Latin America and the Caribbean
 The corresponding references for this table are included in supplementary file 2

The data that were extracted identified nursing registration boards and requirements $n=14$, or professional standards $n=1$.

Table 5 identifies sources of data related to the training and continuing education and development of nurses providing critical care in LICs/LMICs and maps them to regions and countries.


Table 5: Literature providing evidence of training and continuing education available to nurses providing critical care in low income and lower middle-income countries grouped thematically and mapped to region and country

Data	Region	Citations by country
Critical care training courses	SSA	Kenya[1-6], Liberia[7], Malawi[8], Mozambique[9, 10], Rwanda [11, 12],

		Tanzania[13-15], Zambia[16], Multiple countries[17]
	SA	India[18], Nepal[19], Pakistan[20], Sri Lanka[21-24]
	EAP	Myanmar[25], Philippines[26], Vietnam[27]
	MENA	Egypt[28, 29], Palestine[30]
	LAC	Haiti[31]
<i>Education activities targeting specific critical care nursing practices</i>	SSA	Malawi[32], Rwanda[33], Zambia[34]
	SA	India[35-43], Nepal[44], Pakistan[45]
	EAP	Cambodia[46], Myanmar[25], Philippines[26], Vietnam[27], Unnamed EAP country[47]
	MENA	Egypt[29, 48-51], Palestine[30], Multiple MENA countries[52]
	LAC	None
<i>Access to educational resources</i>	SSA	Mozambique[10], Nigeria[53], Tanzania[13], Uganda[54, 55]
	SA	None
	EAP	Philippines[56, 57]
	MENA	Egypt[28, 48, 49, 58]
	LAC	Haiti[59]
<i>Research reporting on before and after testing of educational interventions</i>	SSA	Kenya[3, 60], Malawi[61], Tanzania[62], Uganda[63]
	SA	India[35, 38, 39, 41], Sri Lanka[21]
	EAP	Cambodia[46], Vietnam[27], Unnamed EAP country[47]
	MENA	Egypt[29, 49-51], Palestine[30], Multiple MENA countries[52]
	LAC	None
<i>SSA: Sub Saharan Africa, SA: South Asia, EAP: East Asia and the Pacific, Mena: Middle East and North Africa, LAC: Latin America and the Caribbean</i>		
<i>The corresponding references for this table are included in supplementary file 2</i>		

Data extracted that identified training could be mapped thematically to 1) Education targeting specific practices $n=26$, 2) formal post-graduate study $n=10$, 3) short format courses $n=19$, 4) training courses without a description $n=4$, 5), and critical care training within the undergraduate curriculum $n=3$. Figure 6 maps these themes to region.

Figure 6: Distribution of the literature identifying training and education of nurses providing critical care in LICs/LMICs



The chart shows the distribution of included papers that report on or discuss CCN training and education in LICs/LMICs

Finally, data were extracted identifying access to educational resources including the use of information technologies (IT). Themes included: 1) the availability of peer reviewed literature $n=4$, 2) access to locally generated learning materials or education teams $n=3$, 3) access to IT based education $n=2$, 4) perceived inadequacy of access to educational resources $n=3$.

Discussion

This is the first review to explore and map the evidence related to how the CCN role is performed in LICs/LMICs. Despite the large number of papers meeting inclusion criteria, few provided a comprehensive description of how CCN is performed in any individual LIC/LMICs or region. However, some themes were able to be synthesised from numerous sources and are discussed below.

Who is involved in providing CCN in LICs/LMICs?

Qualified nurses

Documents indicating that registered or qualified nurses were engaged in the care of critically ill patients were found in all regions except LAC. Where details of basic qualifications of CCN's were provided, they included certificate, diploma, and bachelor's degrees with diploma and bachelor's being the most common in many settings. In a large observational study in European hospitals, it was found that for every 10% increase in nurses holding bachelor's degrees rather than a diploma, a 7% reduction in 30-day inpatient mortality was measured.[10] To our knowledge no similar study has been undertaken in resource-limited health systems offering an opportunity for future research.

Nurses with post basic qualification in critical care

The presence of nurses with post-basic training in critical care was reported in all regions but not all countries. Generally, post-basic preparation was reported in LMICs but absent in LICs. Most included literature did not identify the numbers or ratios of critical care trained nurses within each country or region. However, in a Kenyan paper 50% of nurses working in ICUs were reported as holding post-basic qualification.[20] Such a ratio achieves the minimum standards recommended in some HICs, for example, Australia.[11]

Guidelines from India recommend that all nurses working in critical care areas hold formal post-basic qualifications[3] yet even within India this level of preparation is inconsistent.[21] Literature from Sri Lanka and Zambia described situations where the overall percentage of nurses with post-basic qualification was low. However, it was more common that the nurse in charge held a post-basic qualification,[22, 23]. Another situation reported from Zambia suggested, that post-basic trained nurses have responsibilities to provide care to deteriorating patients in the wider hospital reducing the access to trained nurses within the units.[24]

Other cadres of nurses were identified as providing some or all the care commonly associated with the CCN role. Commonly, these included enrolled nurses,[25, 26] nurse-midwives, or

midwife technicians.[27, 28] It was also noted that pre-registration student nurses were engaged in clinical placements within several ICUs and other critical care settings.[27-29]

Non-nursing cadres

Papers from SSA, EAP, and MENA provided evidence of non-nursing staff providing what might be considered aspects of CCN. Roles included technicians[20] and personnel described as clinical officers or non-physician clinicians. However, it was not clear in the included papers what constituted a non-physician clinician[30, 31] Additionally, nurse anaesthetists[32] and respiratory therapists[33] were involved in the care of mechanically ventilated patients. Less clearly defined roles such as healthcare assistants,[23] ancillary staff and caregivers[34-36] were also noted.

Non-healthcare trained carers

A small number of documents suggested that lay-persons provided nursing care. Some of these discussed the parents' roles in neonatal care[37, 38] and were similar to practices in HIC neonatal ICUs such as kangaroo care.[39] Others highlighted the role of lay persons when shortages of nursing staff existed.[22] A global shortage of healthcare workers, predominately in LMIC contexts is well recognised.[9] Urgent work to address this is required if UHC and the attainment of the SDGs[40] are to be met. Although driven by necessity in these examples, multi-directional benefits are reported when families are directly involved in patient care.[41] However, research in critical care environments is sparse, especially in LIC/LMIC settings.[42]

The day-to-day roles and responsibilities of those providing nursing care of critical care patients in LICs/LMICs

Critical care nursing role, practices, or tasks

Although, specific tasks and roles such as care of the ventilated patient, medication management, and clinical education were common, comprehensive description of the role of those providing CCN within the targeted health systems was not well described. However, three papers presented information on required competencies for critical care nurses,[24, 30, 43] thus identifying their broad roles, tasks, and responsibilities. Further documents described the roles of sub-speciality nurses in cardiac-surgical and cardiac-medical units.[44]

CCN systems/staffing/capacity

Details including the numbers of CCNs in the workforce and nursing models of care were common. Two themes in particular stand out: 1) The heterogeneity in the nurse-to-patient ratios reported across regions, countries, and within countries. 2) The perceptions of nurses regarding the conditions they work within.

Reported nurse-to-patient ratios were inconsistent within the included literature. Many reported ratios equivalent to those recommended in many HIC standards for ICUs at 1:1 or 1:2.[11] However, evidence of ratios as low as 1:15 was noted[27] and a perceived inadequacy in the nurse-to-patient ratios to achieve patient care was common. It has been suggested that lower nurse-to-patient ratios is associated with higher mortality rates in ICU [45] although, this is not without controversy.[46] Associations between lower ratios and quality of care[47] and a negative impact on nurse wellbeing[48] have also been reported in HIC contexts.

It remains unclear what would constitute an appropriate nurse-to-patient ratio in resource limited ICUs and warrants further research. The heterogeneity of critical care in these contexts suggests that bespoke recommendations would be required. The use of validated instruments such as the Therapeutic Intervention Scoring System or Nursing Activities Score to assess workload on a unit-based scale may be useful in determining needs. However, even

validated instruments such as these have limitations and do not capture non-clinical tasks or the role of the broader team in providing patient care.[49]

There is evidence that some nurses perceived themselves to be working in poor conditions.[50-52] In the Solomon Islands it was noted that no ICU existed, and nurses undertook all critical care of patients in wards or in non-hospital healthcare settings.[53] Data from a large international survey conducted by the World Federation of Critical Care Nurses found the most common theme, across all regions, was insufficient workforce numbers.[54] This has been highlighted as a priority area of development by the WHO.[9]

In addition to the workforce constraints already discussed, basic commodities including electricity, oxygen, and water may not be guaranteed.[55] Work to identify appropriate technologies to build critical care capacity in resource limited contexts is, therefore, important.[56]

Availability and utilisation of physical resources or technologies to fulfil CCN role

In literature from SSA, SA, and MENA, advanced critical care technologies like those found in high-resource health systems were reported. Examples included mechanical ventilators,[51, 57-60] invasive haemodynamic monitoring[61, 62], and renal replacement therapy.[36, 63] However, it was noted in an Indian paper that the available mechanical ventilation devices, modalities, and adjunctive therapies remained underutilised due to a lack of training on their use.[64] Similarly, where equipment was available, it may be old or poorly functioning in some contexts including Nepal and Sri Lanka.[51, 59]

It should be noted that although critical care is synonymous with high technology resources in HIC's, the associated high costs and need for highly trained staff may not be achievable in resource-limited settings.[8] Research into what constitutes appropriate critical care technologies in resource-limited settings is important.[56] However, critically ill patients

exist in all hospitals, regardless of the availability of equipment and resources. Therefore, delivering critical care should not be predicated on the availability of technology or the ability to achieve all aspects of care in all settings. Instead, critical care should be based on what is appropriate and feasible in each setting.[65]

Availability of policies or guidelines to support evidence-based practice

Limited data spoke to policies, documentation, or practice guidelines in each region. However, a common theme was a perceived absence or inadequacy of these resources. For example in Nigeria, this absence was reported as leading to inadequate knowledge of evidence-based practices.[66] In a Zambian context it was highlighted that even when protocols existed, they were not able to be successfully implemented. However, the reasons for this were not described.[22]. In a review by the same author literature from SSA contexts suggested guidelines were often inappropriate having been developed in HIC's, where different presentations and challenges are experienced.[24]

Conversely, some documents demonstrated the presence of effective policies and practice guidelines, including the generation of locally developed documents. Examples included a comprehensive identification of key documents related to sepsis management in Uganda[67] and a combination of international and locally developed guidelines for care in Nepal.[61]

Registration and professional scope of nurses providing nursing care of critical care patients in LICs/LMICs

Nursing registration and professional standards

Little data were found relating to CCN scope of practice, standards, or registration requirements. Available data from SSA, SA, EAP, and MENA mostly identified the nursing boards that nurses registered under. A small number of papers from SSA offered extended details, such as the continuing professional development requirements for maintaining

registration.[24, 43, 52] Research exploring CCN standards, requirements for registration; including CPD requirements and current scope of practices is needed to guide capacity and health service development in LICs/LMICs.

Training and continuing education available to nurses providing critical care in LICs/LMICs

Formal post-graduate courses

Speciality education in CCN is well documented in many HIC's. For example, in the Australian context a Master's degree program exists in many nursing faculties, with exit points at graduate certificate, post-graduate diploma and master's degree.[68] Where post-basic training was reported in resource-limited settings, only one paper from Zambia provided an overview of course structure or content and provided evidence that the available courses were validated to meet international standards.[24, 69] This lack of detail makes comparison between countries and assessment of the quality of post-basic qualifications difficult. Therefore, presenting an opportunity for research, benchmarking, and curriculum development.

A smaller sub-set of critical care courses were reported where nurses from low-income health systems travelled to other countries to undertake training. One example is found in Rwanda where nurses train in Kenya or South Africa.[70] Caution is required to ensure training attained in this way is contextually appropriate. It was noted by Bruce[71] that CCNs returning to Mozambique following clinical internship in South Africa felt disempowered. Technologies and practices they had experienced were not achievable in their home contexts. It is also evident that healthcare workers from resource-limited countries who are able to train in HIC's frequently stay in those countries once graduated.[72]

It is widely noted that low-wages are both a cause of dissatisfaction and a barrier to nurses pursuing post-basic qualifications in LIC/LMICs.[73] Furthermore, a recent global survey of critical care nursing organisations highlights that credentialling is of great importance to critical care nurses in low income settings.[54] Strategies to make post-basic training accessible and attractive to nurses providing critical care in LICs/LMICs are, therefore, important and an area that deserves attention in research and capacity development agendas.

Short format critical care courses

Many short format courses were developed in-country or developed with support from international collaborators. Some were delivered by local faculty and others by visiting international educators. Among others, the Fundamental Critical Care Support/Paediatric Fundamental Critical Care Support courses in Haiti, Kenya, and Nepal,[30, 57, 74] the Sugar, Temperature, Airway, Blood pressure, Lab work, and Emotional support program,[27, 75] and the Paediatric Assessment of Illness, Recognition and Resuscitation courses[76] were identified in SSA. In SA courses included: The Facility Based Integrated Management of Neonatal and Childhood Illnesses program,[77] the Network for Improving Critical care Skills Training (NICST) program,[78-80] and the Basic Assessment and Support in Intensive Care course.[81, 82]

Educational activities promoted by CCN societies

Tier-3 evidence demonstrated that nursing organisations in SSA, SA, and EAP promoted locally delivered educational workshops or critical care short courses.[83-88] One course in Ghana was promoted as accredited and in partnership with a university nursing faculty.[84] Local and regional conferences were promoted or organised by most of the organisations.[83-90]

Educational activities targeting specific CCN practices

A broad range of educational activities targeting specific CCN practices were noted. Some evidenced informal in-service type of education. However, some nurses in Tanzania and Egypt perceived there to be a gap in this type of job facility-based training.[91, 92] Literature from EAP and SA focussed on task or practice-specific training in areas as diverse as crisis resource management,[93] infection control practices,[94] or the use of physical restraints.[95]

Access to educational resources/Use of IT

Data regarding access to learning materials were extracted. A small number of documents from SSA suggested that access to open-source journals existed,[29] or that learning guides and reading materials were supplied as part of CCN courses.[35, 96] However, papers from Nigeria and Uganda highlighted that a lack of access to relevant evidence and materials was experienced by nurses.[66, 97] Where short-format CCN courses were discussed, it was suggested that the use of low-fidelity simulation[96] and using a train-the-trainer model to upskill clinical nurses in effective clinical teaching were useful strategies.[35] In Tanzania it was suggested that in-service trainings were not a usually practiced.[91] Whereas, in other SSA settings IT capabilities, including limited access to computers[29, 63] were perceived as constraints on the delivery of clinical education.

Educational research

Educational activities targeting specific CCN practices and broader critical care courses were both the subject of educational research in LICs/LMICs. Before and after testing of knowledge,[21] measures of satisfaction and acceptability of training,[77] practice change, and the impact of education on clinical outcomes[98, 99] were all noted. The broader literature suggests that continuing education for nurses in LICs/LMICs remains poorly researched. A recent narrative synthesis identified a range of strategies employed in general

nurse education, not specifically CCN.[100] These included train-the-trainer models, low-dose/high-frequency models, and the use of multiple media; including web-based delivery for training that have been studied in LIC/LMIC contexts. However, they noted the quality of the evidence produced was low.

Great opportunity exists to undertake focussed research investigating the roles, capacity, and continuing education requirements of nurses providing critical care in LICs/LMICs. Research regarding the optimal strategies for delivery of continuing education in these contexts is also needed. Such work is vital in the ongoing development of critical care services in resource constrained health systems.

Addressing capacity development in LICs/LMICs is challenging. Perceived needs may be high but current assets and capacity variable. It is recommended by some that a hybrid approach should be taken where assets, capacity and needs are assessed simultaneously, driven by the local stakeholders.[101] The outcomes of such an assessment should inform contextually appropriate initiatives, that value and leverage current assets and capacity, understand the enablers and barriers to engaging in capacity development work, and clearly identify where needs exist and can be addressed.

Limitations

Although the search strategy was developed to be as comprehensive as possible. Resource constraints, not least the inability to translate non-English language documents, mean it is possible that some literature was not captured. Efforts to access the websites of national health departments and nursing councils or directly contact these organisations may elucidate registration requirements, scopes, and standards of practice. However, it remains unclear whether this would provide data specific to CCN. Finally, the lack of literature found in

English from LICs/LMICs in LAC prevents an accurate picture of CCN in these contexts to be drawn.

Conclusion

Provision of critical care is a complex undertaking and despite commonalities, is unique to regional and socio-economic contexts. Multi-disciplinary teamwork is paramount. Yet the structure and skill levels of such teams will vary according to patient population, resources, and treatments available. The findings of this study documented care of critically ill patients in specialist and non-specialist environments. Human and physical resourcing was highly variable across regions but often reported as inadequate to meet demand within LIC/LMIC health systems. The descriptions of CCN training and skill levels were equally variable but often perceived as requiring development. Therefore, a universal definition of the CCN role in LIC/LMIC health systems is likely unhelpful. Research opportunities abound to elucidate current assets, capacity and needs of CCNs in specific LIC/LMIC contexts. Outputs from such research would be invaluable in supporting contextually appropriate capacity development programs.

Funding Statement

This research is supported by an Australian Government Research Training Program (RTP) Scholarship. Award/Grant number is not applicable.

Contributorship statement

Made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data: AM, GOR, GW, PC.

Involved in drafting the manuscript or revising it critically for important intellectual content: AM, GOR, GW, PC.

Given final approval of the version to be published. Each author should have participated sufficiently in the work to take public responsibility for appropriate portions of the content:

AM, GOR, GW, PC.

Agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved: AM,

GOR, GW, PC.

Conflict of interest statement

No conflict of interest has been declared by the authors.

Data availability statement

No additional data available.

Figure captions

PREPRINT

References

1. Marshall JC, Bosco L, Adhikari NK, Connolly B, Diaz JV, Dorman T, et al. What is an intensive care unit? A report of the task force of the World Federation of Societies of Intensive and Critical Care Medicine. *Journal of Critical Care*. 2017;37:270-6.
2. Murthy S, Leligdowicz A, Adhikari NKJ. Intensive Care Unit Capacity in Low-Income Countries: A Systematic Review. *PLoS ONE*. 2015;10(1):1-12.
3. Rungta N, Zirpe KG, Dixit SB, Mehta Y, Chuadhry D, Govil D, et al. Indian Society of Critical Care Medicine Experts Committee Consensus Statement on ICU Planning and Designing, 2020. *Indian Journal of Critical Care Medicine*. 2020;24:S43-S60.
4. Adhikari NKJ, Fowler RA, Bhagwanjee S, Rubenfeld GD. Critical care and the global burden of critical illness in adults. *The Lancet*. 2010;376(9749):1339-46.
5. Vincent J-L, Marshall JC, Namendys-Silva SA, François B, Martin-Loeches I, Lipman J, et al. Assessment of the worldwide burden of critical illness: the Intensive Care Over Nations (ICON) audit. *The Lancet Respiratory Medicine*. 2014;2(5):380-6.
6. Vukoja M, Riviello ED, Schultz MJ. Critical care outcomes in resource-limited settings. 2018. p. 421-7.
7. Vukoja M, Riviello E, Gavrilovic S, Adhikari NKJ, Kashyap R, Bhagwanjee S, et al. A Survey on Critical Care Resources and Practices in Low- and Middle-Income Countries. *Global Heart*. 2014;9(3):337-42.
8. Diaz JV, Riviello ED, Papali A, Adhikari NKJ, Ferreira JC. Global Critical Care: Moving Forward in Resource-Limited Settings. 2019.
9. WHO. Global strategy on human resources for health: workforce 2030. Geneva, Switzerland: World Health Organization; 2016.
10. Aiken LHP, Sloane DMP, Bruyneel LMS, Van den Heede KP, Griffiths PP, Busse RP, et al. Nurse staffing and education and hospital mortality in nine European countries: a retrospective observational study. *Lancet*. 2014;383(9931):1824-30.
11. Chamberlain D, Pollock W, Fulbrook P. ACCCN Workforce Standards for Intensive Care Nursing: Systematic and evidence review, development, and appraisal. *Australian Critical Care*. 2018(5).
12. Bray K, Wren I, Baldwin A, St Ledger U, Gibson V, Goodman S, et al. Standards for nurse staffing in critical care units determined by: The British Association of Critical Care Nurses, The Critical Care Networks National Nurse Leads, Royal College of Nursing Critical Care and In-flight Forum. *Nurs Crit Care*. 2010;15(3):109-11.
13. Chamberlain D, Pollock W, Fulbrook P. ACCCN Workforce Standards for Intensive Care Nursing: Systematic and evidence review, development, and appraisal. *Aust Crit Care*. 2018;31(5):292-302.
14. Peters MDJ, Godfrey C, McInerney P, Munn Z, Tricco AC, Khalil H. Scoping reviews. In: Aromataris E MZ, editor. *JBIM Manual for Evidence Synthesis*: JBI; 2020.
15. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Annals Of Internal Medicine*. 2018;169(7):467-73.
16. Macey A. The critical care nursing role in low and lower-middle income settings: A scoping review protocol 2020.
17. Pham MT, Rajić A, Greig JD, Sargeant JM, Papadopoulos A, McEwen SA. A scoping review of scoping reviews: advancing the approach and enhancing the consistency. *Res Synth Methods*. 2014;5(4):371-85.
18. Kepes S, Banks GC, McDaniel M, Whetzel DL. Publication Bias in the Organizational Sciences. *Organizational Research Methods*. 2012;15(4):624-62.
19. The World Bank Group. The world by region 2018 [Available from: <https://datatopics.worldbank.org/sdgatlas/archive/2018/the-world-by-region.html>].

20. Waweru-Siika W, Mung'ayi V, Misango D, Mogi A, Kisia A, Ngumi Z. The history of critical care in Kenya. *J Crit Care*. 2020;55:122-7.
21. Dongara AR, Nimbalkar SM, Phatak AG, Patel DV, Nimbalkar AS. An Educational Intervention to Improve Nurses' Understanding of Pain in Children in Western India. *Pain Manag Nurs*. 2017;18(1):24-32.
22. Carter C, Snell D. Nursing the critically ill surgical patient in Zambia. *Br J Nurs*. 2016;25(20):1123-8.
23. Haniffa R, De Silva AP, Iddagoda S, Batawalage H, De Silva S, Mahipala PG, et al. A cross-sectional survey of critical care services in Sri Lanka: A lower middle-income country. *Journal of Critical Care*. 2014;29(5):764-8.
24. Carter C, Mukonka PS, Sitwala LJ, Howard-Hunt B, Notter J. The development of critical care nursing education in Zambia. *British journal of nursing (Mark Allen Publishing)*. 2020;29(9):499-505.
25. Ufashingabire CM, Nsereko E, Njunwa KJ, Brysiewicz P. Knowledge and attitudes of nurses regarding pain in the intensive care unit patients in Rwanda. *Rwanda Journal Series F: Medicine and Health Sciences*. 2016;3(1):21-6.
26. Mwewa B, Mweemba P. Knowledge and utilization of ICU admission criteria and guidelines, Lusaka, Zambia. *Medical Journal of Zambia*. 2010;37(3):143-52.
27. Guiles S, Lemons J, Trautman M, Bucher S, Songok J, Gisore P. The Implementation of a Neonatal Nurse Training Program at the Riley Mother Baby Hospital of Kenya. *Newborn Infant Nurs Rev*. 2016;16(4):184-9.
28. Tadyanemhandu C, van Aswegen H, Ntsiea V. Early mobilisation practices of patients in intensive care units in Zimbabwean government hospitals - a cross-sectional study. *South Afr J Crit Care*. 2018;34(1):16-21.
29. Bruce JC, Dippenaar J, Schmollgruber S, Mphuthi DD, Huiskamp A. Advancing nursing scholarship: the Mozambique model. *Global Health Action*. 2017;10:1351116.
30. Macleod JBA, Jones T, Aphivantrakul P, Chupp M, Poenaru D. Evaluation of fundamental critical care course in Kenya: knowledge, attitude, and practice. *The Journal of surgical research*. 2011;167(2):223-30.
31. Baker T, Lugazia E, Eriksen J, Mwafongo V, Irestedt L, Konrad D. Emergency and critical care services in Tanzania: a survey of ten hospitals. *Bmc Health Services Research*. 2013;13:9.
32. Okech U, Chokwe T, Mung'ayi V. The operational setup of intensive care units in a low income country in East Africa. *East African medical journal*. 2015;92(2):72-80.
33. Sawe HR, Mfinanga JA, Lidenge SJ, Mpondo BC, Msangi S, Lugazia E, et al. Disease patterns and clinical outcomes of patients admitted in intensive care units of tertiary referral hospitals of Tanzania. *BMC Int Health Hum Rights*. 2014;14(1):26.
34. Spiwak R, Lett R, Rwanyuma L, Logsetty S. Creation of a standardized burn course for Low Income Countries: Meeting local needs. *Burns*. 2014;40(7):1292-9.
35. Cunningham C, Brysiewicz P, Sepeku A, White L, Murray B, Lobue N, et al. Developing an emergency nursing short course in Tanzania. *Afr J Emerg Med*. 2017;7(4):147-50.
36. Atumanya P, Sendagire C, Wabule A, Mukisa J, Ssemogerere L, Kwizera A, et al. Assessment of the current capacity of intensive care units in Uganda; A descriptive study. *Journal of Critical Care*. 2020;55:95-9.
37. Sathish Y, Lewis LE, Noronha JA, Nayak BS, Pai MS, Altimier L. Promoting developmental supportive care in preterm infants and families in a level III neonatal intensive care unit (NICU) setting in India. *Nurse Education in Practice*. 2019;40:8.
38. Phuma-Ngaiyaye E, Kalembo FW. Supporting mothers to bond with their newborn babies: Strategies used in a neonatal intensive care unit at a tertiary hospital in Malawi. *International Journal of Nursing Sciences*. 2016;3(4):362-6.
39. Altimier L, Phillips RM. The Neonatal Integrative Developmental Care Model: Seven Neuroprotective Core Measures for Family-Centered Developmental Care. *Newborn Infant Nurs Rev*. 2013;13(1):9-22.

40. United Nations. Sustainable Development Goals 2021 [Available from: <https://sdgs.un.org/goals/goal3>].
41. Jazieh AR, Volker S, Taher S. Involving the Family in Patient Care: A Culturally Tailored Communication Model. *Global Journal on Quality and Safety in Healthcare*. 2020;1(2):33-7.
42. Olding M, McMillan SE, Reeves S, Schmitt MH, Puntillo K, Kitto S. Patient and family involvement in adult critical and intensive care settings: a scoping review. *Health Expect*. 2016;19(6):1183-202.
43. Gundo R. Development, Implementation and Evaluation of an In-service Training Programme for Critical Care Nurses in Malawi: Auckland University of Technology; 2019.
44. Dongara AR, Shah SN, Nimbalkar SM, Phatak AG, Nimbalkar AS. Knowledge of and Attitudes Regarding Postoperative Pain among the Pediatric Cardiac Nursing Staff: An Indian Experience. *Pain Manag Nurs*. 2015;16(3):314-20.
45. Driscoll A, Grant MJ, Carroll D, Dalton S, Deaton C, Jones I, et al. The effect of nurse-to-patient ratios on nurse-sensitive patient outcomes in acute specialist units: a systematic review and meta-analysis. *Eur J Cardiovasc Nurs*. 2018;17(1):6-22.
46. Margadant C, Wortel S, Hoogendoorn M, Bosman R, Spijkstra JJ, Brinkman S, et al. The Nursing Activities Score Per Nurse Ratio Is Associated With In-Hospital Mortality, Whereas the Patients Per Nurse Ratio Is Not. *Crit Care Med*. 2020;48(1):3-9.
47. Falk AC, Wallin EM. Quality of patient care in the critical care unit in relation to nurse patient ratio: A descriptive study. *Intensive Crit Care Nurs*. 2016;35:74-9.
48. Carayon P, Alvarado CJ. Workload and patient safety among critical care nurse. *Crit Care Nurs Clin North Am*. 2007;19:121-9.
49. Scruth E. Nursing Activities Score, Nurse Patient Ratios, and ICU Mortality: Its More Complicated Than That. *Critical care medicine*. 2020;48(1):126-7.
50. Paguio JT, Banayat AC. Commentary on challenges to critical care nursing practice in the Philippines. *The World of Critical Care Nursing*. 2018;12(1).
51. Limbu S, Kongsuwan W, Yodchai K. Lived experiences of intensive care nurses in caring for critically ill patients. *Nursing in Critical Care*. 2019;24(1):9-14.
52. Gundo R. Contextual issues that influence preparedness of nurses for critical care nursing practice in Malawi. *Malawi Med J*. 2019;31(2):138-43.
53. Westcott M, Martiniuk ALC, Fowler RA, Adhikari NKJ, Dalipanda T. Critical care resources in the Solomon Islands: a cross-sectional survey. *BMC Int Health Hum Rights*. 2012;12:9.
54. Williams G, Fulbrook P, Kleinpell R, Alberto L. The Fifth International Survey of Critical Care Nursing Organizations: Implications for Policy. *J Nurs Scholarsh*. 2020;52(6):652-60.
55. Dondorp A, Iyer S, Schultz M. Critical Care in Resource-Restricted Settings. *JAMA*. 2016(8):753.
56. Turner HC, Nguyen Van H, Yacoub S, Van Minh Tu H, Clifton DA, Thwaites GE, et al. Achieving affordable critical care in low-income and middle-income countries. 2019.
57. Kumar R, Canarie MF. Developing Pediatric Critical Care in Kenya*. *Pediatric Critical Care Medicine*. 2019;20(12):E538-E45.
58. Sawe HR, Mfinanga JA, Lidenge SJ, Mpondo BCT, Msangi S, Lugazia E, et al. Disease patterns and clinical outcomes of patients admitted in intensive care units of tertiary referral hospitals of Tanzania. *BMC Int Health Hum Rights*. 2014;14.
59. Charlton S, O'Reilly G, Jones T, Fitzgerald M. Emergency care in developing nations: The role of emergency nurses in Galle, Sri Lanka. *Australas Emerg Nurs J*. 2011;14(2):69-74.
60. Aloush SM, Abdelkader FA, Al-Sayaghi K, Tawalbeh LI, Suliman M, Al Bashtawy M, et al. Compliance of Nurses and Hospitals With Ventilator-Associated Pneumonia Prevention Guidelines: A Middle Eastern Survey. *Journal of Nursing Care Quality*. 2018;33(3):E8-E14.
61. Khanal A, Sharma A, Basnet S. Current State of Pediatric Intensive Care and High Dependency Care in Nepal. *Pediatric Critical Care Medicine*. 2016;17(11):1032-40.

62. Baker T, Lugazia E, Eriksen J, Mwafongo V, Irestedt L, Konrad D. Emergency and critical care services in Tanzania: a survey of ten hospitals. *BMC Health Services Research*. 2013;13.
63. Marzuola JM. Teaching Strategies to Reduce VAP at Mulago Hospital: A Capstone Project. 2016.
64. Atreya MR, Lorenz JM, Narendran V. Provider Perceptions of Bubble Continuous Positive Airway Pressure and Barriers to Implementation in a Level III Neonatal Unit in South India. *Adv Neonatal Care*. 2018;18(6):500-6.
65. Riviello ED, Letchford S, Achieng L, Newton MW. Critical care in resource-poor settings: Lessons learned and future directions. *Crit Care Med*. 2011;39(4):860-7.
66. Ilesanmi RE, Olabisi P. Assessment of Common Interventions and Perceived Barriers to Pressure Ulcer Prevention in Southwest Nigeria. *J Wound Ostomy Cont Nurs*. 2014;41(3):242-6.
67. Nakate GM, Emelonye AU. Examining nursing practices for management of sepsis in low income countries: the case of Uganda. *World*. 2018;12(1):2.
68. Gill FJ, Leslie GD, Grech C, Latour JM. An analysis of Australian graduate critical care nurse education. *Collegian*. 2015;22(1):71-81.
69. UNESCO. International standard classification of education. UNESCO Institute of statistics; 2011.
70. Munyiginya P, Brysiewicz P, Mill J. Critical care nursing practice and education in Rwanda. *South Afr J Crit Care*. 2016;32(2):55-7.
71. Bruce JC, Schmollgruber S, Baumann J. Intercountry master's degree in nursing: policy implications for the Mozambican health system. *International Nursing Review*. 2018;65(3):425-33.
72. OECD. Recent Trends in International Migration of Doctors, Nurses and Medical Students 2019.
73. Ng'ang'a N, Byrne MW. Professional practice models for nurses in low-income countries: an integrative review. *BMC Nurs*. 2015;14(1):44-.
74. Hernandez J, Kioko M, Adeyinka A, Pinto R, Mercredi G, Petrescu M, et al. Pediatric fundamentals of critical care support (PFCCS) course performance among staff from U.S. and resource-limited countries. *Pediatric Critical Care Medicine*. 2012;13(6):718.
75. Kendall AB, Scott PA, Karlsen KA. The S.T.A.B.L.E.[®] Program: The Evidence Behind the 2012 Update. *J Perinat Neonatal Nurs*. 2012;26(2):147-57.
76. MacDonell R, Koe S, Okafor I, James A, Ahern C, Scanlan T. Building resuscitation capability in low resource settings; development of the PAIRS course (outlining the development of a tailored paediatric resuscitation programme for low resource healthcare settings). *Archives of Disease in Childhood*. 2019;104(Supplement 3):A259.
77. Patel DV, Nimbalkar SM, Singh US, Kharod NM. Acceptability and implementation of FIMNCI by medical officers and staff nurses in government health institutions of Western India. *Archives of Disease in Childhood*. 2012;97(SUPPL. 2):A489-A90.
78. De Silva AP, Stephens T, Welch J, Sigera C, De Alwis S, Athapattu P, et al. Nursing intensive care skills training: A nurse led, short, structured, and practical training program, developed and tested in a resource-limited setting. *Journal of Critical Care*. 2015;30(2):438.e7-.e11.
79. Stephens T, De Silva AP, Beane A, Welch J, Sigera C, De Alwis S, et al. Capacity building for critical care training delivery: Development and evaluation of the Network for Improving Critical care Skills Training (NICST) programme in Sri Lanka. *Intensive & critical care nursing*. 2017;39:28-36.
80. Beane A, Stephens T, De Silva A, Adikaram M, De Alwis S, Athapattu P, et al. A collaborative approach to training ward nurses in acute care skills in resource limited settings: the nursing intensive care skills training (nicts) project. *Intensive Care Medicine Experimental*. 2015;3:1-2.
81. Gomersall. Basic assessment and support in intensive care 2018 [Available from: <https://www.aic.cuhk.edu.hk/web8/BASIC.htm>].
82. Gautam P, Acharya SP, Williams G. Formation of the Southern Asia Association for Regional Cooperation Federation of Critical Care Nurses. *The World of Critical Care Nursing*. 2018;12(1).
83. WFCCN. The world federation of critical care nurses 2021 [Available from: <http://wfccn.org>].

84. CCNGG. Critical Care Nurses Group Ghana - CCNGG Facebook site 2020 [
85. CCNAU. Critical Care Nursing Association of Uganda - CCNAU facebook page 2020 [Available from: <https://www.facebook.com/CCNAU.ORG/>.
86. CCNAPI. Critical care nurses association of the philippines website 2020 [Available from: <http://www.ccnapi.org/>.
87. SSCCN. The Sri Lankan society of critical care nurses website 2020 [Available from: <https://www.scccn.org/>.
88. CCNAN. Critical Care Nurses Association of Nepal - CCNAN facebook page 2020 [Available from: <https://www.facebook.com/CCNANNepal/>.
89. AFCCN. Conferences and events 2021 [Available from: <https://www.africanfccn.org/conferences-events>.
90. Nigeria Nw. Nursing world Nigeria 2020 [Available from: <https://www.nursingworldnigeria.com/index.php>.
91. Mbwele B, Ide NL, Mrema JG, Sarah APW, Melnick JA, Manongi R. Learning from Health Care Workers' Opinions for Improving Quality of neonatal Health Care in Kilimanjaro region, northeast Tanzania. *Ann Med Health Sci Res.* 2014;4(1):105-14.
92. Eltaybani S, Abdelwareth M, Abou-Zeid NA, Ahmed N. Recommendations to prevent nursing errors: Content analysis of semi-structured interviews with intensive care unit nurses in a developing country. *Journal of Nursing Management.* 2020;28(3):690-8.
93. Emani SS, Allan CK, Forster T, Fisk AC, Lagrasta C, Zheleva B, et al. Simulation training improves team dynamics and performance in a low-resource cardiac intensive care unit. *Ann Pediatr Cardiol.* 2018;11(2):130-6.
94. Sharma R, Jagota R, Koushal V. Study of Knowledge, Awareness and Practices of Infection Control among ICU Staff of a Multispecialty Tertiary Level Teaching Institute of North India. *J Health Manag.* 2013;15(1):45-56.
95. Pradhan N, Lama S, Mandal G, Shrestha E. Physical restraining: Nurses knowledge and practice in Tertiary Care Hospital of Eastern Nepal. *Nurs Open.* 2019;6(3):1029-37.
96. Byrne-Davis L, Slattery H, Whiteside S, Moro E, Jackson M, Onyachi N, et al. Efficacy and acceptability of an acute illness management course delivered to staff and students in Uganda by staff from the UK. *Int Health.* 2015;7(5):360-6.
97. Kizza IB, Muliira JK. Nurses' pain assessment practices with critically ill adult patients. *International Nursing Review.* 2015;62(4):573-82.
98. Acharya R, Mishra SB, Ipsita S, Azim A. Impact of Nursing Education on CLABSI Rates: An Experience from a Tertiary Care Hospital in Eastern India. *Indian Journal of Critical Care Medicine.* 2019;23(7):316-9.
99. Sahni N, Biswal M, Gandhi K, Kaur K, Saini V, Yaddanapudi LN. Effect of Intensive Education and Training of Nurses on Ventilator-associated Pneumonia and Central Line-associated Bloodstream Infection Incidence in Intensive Care Unit at a Tertiary Care Center in North India. *Indian Journal of Critical Care Medicine.* 2017;21(11):779-82.
100. Azad A, Min J-G, Syed S, Anderson S. Continued nursing education in low-income and middle-income countries: a narrative synthesis. *BMJ Glob Health.* 2020;5(2):e001981.
101. Altschuld J, Hsin-Ling S, Lee Y-F. Needs assessment and asset/capacity building: A promising development in practice. In: James W. Altschuld, Ryan Watkins, editors. *Needs assessment: Trends and a view toward the future: Wiley periodicals; 2014.*

Supplement 1: Search strings, additional limits, and database versions

Database versions accessed

- 1) Ovid MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Daily and Versions(R)
- 2) Embase Classic+Embase
- 3) Ovid Emcare
- 4) Global Health
- 5) SCOPUS
- 6) Web of Science.

Applied search limits

Limits set in all database searches were publications in the last 10 years (2010-2020) and publications in English language.

Additional limits in SCOPUS were imposed to results from 3 subject areas:

- 1) medical
- 2) nursing
- 3) health professions.

Additional limits in Web of science were the categories:

- 1) nursing
- 2) critical care medicine
- 3) education
- 4) educational research
- 5) education scientific disciplines
- 6) health care sciences services
- 7) health policy services

Ovid search strings

((nurs* or nurs* assistant* or Physician* assistant* or Multidisciplinary team* or Care giver* or Staff* or technician*) and (Intensive care or critical care or coronary care)).mp.

((((Role* or practice* or skill* or task* or Responsibilit* or ratio* or Scope of practice* or (Nurs* or profession*)) adj2 patient* relationship*) or education* or Prepar* or train* or development or (intensive care or critical care)) adj3 course*).mp

(((((Low or Develop*) adj3 (countr* or nation*)) or (limit* or restrict or poor or low)) adj3 resource*) or Third world or africa or maghreb or maghrib or sahara or algeria or egypt or morocco or tunisia or "africa south of the sahara" or sub-saharan or subsaharan or cameroon or central african republic or chad or comoros or congo or "democratic republic of the congo" or madagascar or "sao tome and principe" or british indian ocean territory or burundi or djibouti or eritrea or ethiopia or kenya or rwanda or somalia or south sudan or sudan or tanzania or uganda or angola or eswatini or lesotho or malawi or mozambique or zambia or zimbabwe or benin or burkina faso or cabo verde or cote

	<p>d'ivoire or gambia or ghana or guinea or guinea-bissau or liberia or mali or mauritania or niger or nigeria or senegal or sierra leone or togo or haiti or nicaragua or honduras or el salvador or Bolivia or kyrgyzstan or tajikistan or uzbekistan or kiribati or cambodia or laos or myanmar or philippines or timor-leste or vietnam or bangladesh or bhutan or india or nepal or pakistan or sri lanka or "democratic people's republic of korea" or Mongolia or west bank or gaza or afghanistan or syria or yemen or bosnia or moldova or Ukraine or pacific islands or papua new guinea or vanuatu or micronesia or solomon islands).mp.</p>
--	---

PRE-PRIMA

Supplement 2: Bibliographies for tables 2-5

Bibliography of citations in Table 2

1. Kassew T, Tilahun A. D, Liyew B. Nurses' Knowledge, Attitude, and Influencing Factors regarding Physical Restraint Use in the Intensive Care Unit: A Multicenter Cross-Sectional Study. *Crit Care Res Pract.* 2020.
2. Okech UK, Chokwe T, Mung'ayi V. The operational setup of intensive care units in a low income country in East Africa. *EAST AFRICAN MEDICAL JOURNAL.* 2015;92(2).
3. Muniz SA, Lang RW, Falcon L, Garces-King J, Willard S, Peck GL. Preparing Global Trauma Nurses for Leadership Roles in Global Trauma Systems. *J Trauma Nurs.* 2017;24(5):306-11.
4. Gundo R. Contextual issues that influence preparedness of nurses for critical care nursing practice in Malawi. *Malawi Med J.* 2019;31(2):138-43.
5. Gundo R. Development, Implementation and Evaluation of an In-service Training Programme for Critical Care Nurses in Malawi. 2019.
6. Ufashingabire CM, Nsereko E, Njunwa KJ, Brysiewicz P. Knowledge and attitudes of nurses regarding pain in the intensive care unit patients in Rwanda. *Rwanda Journal.* 2016;3(1):21-6.
7. Cunningham C, Brysiewicz P, Sepeku A, White L, Murray B, Lobue N, et al. Developing an emergency nursing short course in Tanzania. *Afr J Emerg Med.* 2017;7(4):147-50.
8. Kizza IB, Muliira JK. Nurses' pain assessment practices with critically ill adult patients. *International Nursing Review.* 2015;62(4):573-82.
9. Carter C, Mukonka PS, Sitwala LJ, Howard-Hunt B, Notter J. The development of critical care nursing education in Zambia. *British journal of nursing (Mark Allen Publishing).* 2020;29(9):499-505.
10. Sathish Y, Lewis LE, Noronha JA, Nayak BS, Pai MS, Altimier L. Promoting developmental supportive care in preterm infants and families in a level III neonatal intensive care unit (NICU) setting in India. *Nurse Education in Practice.* 2019;40:8.
11. Dongara AR, Nimbalkar SM, Phatak AG, Patel DV, Nimbalkar AS. An Educational Intervention to Improve Nurses' Understanding of Pain in Children in Western India. *Pain Manag Nurs.* 2017;18(1):24-32.
12. Kalyan G, Bibi R, Kaur R, Bhatti R, Kumari R, Rana R, et al. Knowledge and practices of intensive care unit nurses related to prevention of ventilator associated pneumonia in selected intensive care units of a tertiary care centre, India. *Iranian Journal of Nursing and Midwifery Research.* 2020;25(5):369-75.
13. Pradhan N, Lama S, Mandal G, Shrestha E. Physical restraining: Nurses knowledge and practice in Tertiary Care Hospital of Eastern Nepal. *Nurs Open.* 2019;6(3):1029-37.
14. Bongar MVV, Pangan FC, Macindo JRB. Characteristics of a critical care clinical placement programme affecting critical care nursing competency of baccalaureate nursing students: A structural equation modelling. *Journal of Clinical Nursing.* 2019;28(9-10):1760-70.
15. Paguio JT, Banayat AC. Commentary on challenges to critical care nursing practice in the Philippines. *The World of Critical Care Nursing.* 2018;12(1).
16. Martinez RCKP, Rogado MIC, Serondo DJF, Soriano GP, Ilano KCS. Critical Care Nursing in the Philippines: Historical Past, Current Practices, and Future Directions. *Critical care nursing clinics of North America.* 2021;33(1):75-87.
17. Westcott M, Martiniuk ALC, Fowler RA, Adhikari NKJ, Dalipanda T. Critical care resources in the Solomon Islands: a cross-sectional survey. *BMC Int Health Hum Rights.* 2012;12:9.
18. Fadadu PP, Liu JC, Schiltz BM, Xoay TD, Phuc PH, Kumbamu A, et al. A Mixed-Methods Exploration of Pediatric Intensivists' Attitudes toward End-of-Life Care in Vietnam. *J Palliat Med.* 2019;22(8):885-93.
19. AL-Gabri MM, Mohammed MA, Mehany MM. Factors Affecting Nurses Compliance Regarding the Care of Patients with Chest Trauma.

20. Al-Rafay SS, Al-Sharkawy SS. Educational Outcomes Associated With Providing a Comprehensive Guidelines Program About Nursing Care of Preterm Neonates Receiving Total Parenteral Nutrition. *Clin Nurs Res.* 2012;21(2):142-58.
21. Attia MAS, Youseff MRL, Abd El Fatah SAM, Ibrahim SK, Gomaa NAS. The relationship between health care providers' perceived work climate, organizational commitment, and caring efficacy at pediatric intensive care units, Cairo University. *Int J Health Plan Manag.* 2020;35(2):469-81.
22. Mohamed SS, Ali BAEM, Abd-El-Karim HE, Masoed ES. Nurses' knowledge and performance regarding infection control during milk formula preparation and its effect on neonates at El-Minia city hospitals. *The Journal of American Science.* 2012;8(10):782-7.
23. Selim AA, Ely EW. Delirium the under-recognised syndrome: survey of healthcare professionals' awareness and practice in the intensive care units. *Journal of Clinical Nursing.* 2017;26(5-6):813-24.
24. Eltaybani S, Abdelwareth M, Abou-Zeid NA, Ahmed N. Recommendations to prevent nursing errors: Content analysis of semi-structured interviews with intensive care unit nurses in a developing country. *Journal of Nursing Management.* 2020;28(3):690-8.
25. Aloush SM, Abdelkader FA, Al-Sayaghi K, Tawalbeh LI, Suliman M, Al Bashtawy M, et al. Compliance of Nurses and Hospitals With Ventilator-Associated Pneumonia Prevention Guidelines: A Middle Eastern Survey. *Journal of Nursing Care Quality.* 2018;33(3):E8-E14.
26. Gorman LL, McDowell JRS. Identifying the needs of critical and acute cardiac care nurses within the first two years of practice in Egypt using a nominal group technique. *Nurse Education in Practice.* 2018;28:127-34.
27. Khallil NS. Critical care nurses' use of non-pharmacological pain management methods in Egypt. *Appl Nurs Res.* 2018;44:33-8.
28. Kandeel N, Ibrahim Y. Student Nurses' Perception on the Impact of Information Technology on Teaching and Learning. *Int J Inf Commun Technol Educ.* 2010;6(3):38-50.
29. Zyoud SH, Khaled SM, Kawasmi BM, Habeba AM, Hamadneh AT, Anabosi HH, et al. Knowledge about the administration and regulation of high alert medications among nurses in Palestine: a cross-sectional study. *Bmc Nursing.* 2019;18.
30. Opare MA. Ghana. In: Klopper HC, Uys LR, editors. *The State of Nursing and Nursing Education in Africa: A Country-by-Country Review.* Indianapolis: Sigma Theta Tau International; 2013. p. 93-112.
31. Kimani G, Maina F, Rithara S. Perception of palliative care course among other programs in a government learning institution. *Supportive Care in Cancer.* 2017;25(2 Supplement 1):S106.
32. Waweru-Siika W, Mung'ayi V, Misango D, Mogi A, Kisia A, Ngumi Z. The history of critical care in Kenya. *J Crit Care.* 2020;55:122-7.
33. Karani AK, Milimo BW, Kuria RW. Kenya. In: Klopper H, Uys L, editors. *The State of Nursing and Nursing Education in Africa: A Country-by-Country Review.* Indianapolis: Sigma Theta Tau International; 2013. p. 113-26.
34. Bruce JC, Schmollgruber S, Baumann J. Intercountry master's degree in nursing: policy implications for the Mozambican health system. *International Nursing Review.* 2018;65(3):425-33.
35. Mukamana D. Rwanda. In: Klopper H, Uys LR, editors. *The State of Nursing and Nursing Education in Africa: A Country-by-Country Review.* Indianapolis: Sigma Theta Tau International; 2013. p. 231-47.
36. Sawe HR, Mfinanga JA, Lidenge SJ, Mpondo BCT, Msangi S, Lugazia E, et al. Disease patterns and clinical outcomes of patients admitted in intensive care units of tertiary referral hospitals of Tanzania. *BMC Int Health Hum Rights.* 2014;14.
37. Baker T, Lugazia E, Eriksen J, Mwafongo V, Irestedt L, Konrad D. Emergency and critical care services in Tanzania: a survey of ten hospitals. *BMC Health Services Research.* 2013;13.

38. Atumanya P, Sendagire C, Wabule A, Mukisa J, Ssemogerere L, Kwizera A, et al. Assessment of the current capacity of intensive care units in Uganda; A descriptive study. *Journal of Critical Care*. 2020;55:95-9.
39. Carter C, Snell D. Nursing the critically ill surgical patient in Zambia. *Br J Nurs*. 2016;25(20):1123-8.
40. Tadyanemhandu C, van Aswegen H, Ntsiea V. Early mobilisation practices of patients in intensive care units in Zimbabwean government hospitals - a cross-sectional study. *South Afr J Crit Care*. 2018;34(1):16-21.
41. Rungta N, Zirpe KG, Dixit SB, Mehta Y, Chuadhry D, Govil D, et al. Indian Society of Critical Care Medicine Experts Committee Consensus Statement on ICU Planning and Designing, 2020. *Indian Journal of Critical Care Medicine*. 2020;24:S43-S60.
42. Haniffa R, De Silva AP, Iddagoda S, Batawalage H, De Silva S, Mahipala PG, et al. A cross-sectional survey of critical care services in Sri Lanka: A lower middle-income country. *Journal of Critical Care*. 2014;29(5):764-8.
43. Stephens T, De Silva AP, Beane A, Welch J, Sigera C, De Alwis S, et al. Capacity building for critical care training delivery: Development and evaluation of the Network for Improving Critical care Skills Training (NICST) programme in Sri Lanka. *Intensive & critical care nursing*. 2017;39:28-36.
44. Charlton S, O'Reilly G, Jones T, Fitzgerald M. Emergency care in developing nations: The role of emergency nurses in Galle, Sri Lanka. *Australas Emerg Nurs J*. 2011;14(2):69-74.
45. Barkley AS, Medina-Beckwith J, Sothea S, Pak S, Durfy SJ, Lele AV. Teaching the Emergency Neurologic Life Support Course at Two Major Hospitals in Phnom Penh, Cambodia. *World neurosurgery*. 2020.
46. Losonczy LI, Barnes SL, Liu S, Williams SR, McCurdy MT, Lemos V, et al. Critical care capacity in Haiti: A nationwide cross-sectional survey. *PLoS One*. 2019;14(6):e0218141.
47. Guiles S, Lemons J, Trautman M, Bucher S, Songok J, Gisore P. The Implementation of a Neonatal Nurse Training Program at the Riley Mother Baby Hospital of Kenya. *Newborn Infant Nurs Rev*. 2016;16(4):184-9.
48. Phuma-Ngaiyaye E, Kalembo FW. Supporting mothers to bond with their newborn babies: Strategies used in a neonatal intensive care unit at a tertiary hospital in Malawi. *International Journal of Nursing Sciences*. 2016;3(4):362-6.
49. Bruce JC, Dippenaar J, Schmollgruber S, Mphuthi DD, Huiskamp A. Advancing nursing scholarship: the Mozambique model. *Global Health Action*. 2017;10:1351116.
50. Macindo JRB, Danganan MPB, Soriano CAF, Kho NSR, Bongar MVV. A conjoint analysis of the acute and critical care experiential learning preferences of Baccalaureate student nurses. *Nurse Education in Practice*. 2019;36:125-31.
51. Zahran EM, Taha EE. Integrating evidence based nursing into the critical care nursing course: challenges from students' perspectives. *The Journal of American Science*. 2011;7(7):135-44.
52. MacLeod JBA, Jones T, Aphivantrakul P, Chupp M, Poenaru D. Evaluation of fundamental critical care course in kenya: Knowledge, attitude, and practice. *Journal of Surgical Research*. 2011;167(2):223-30.
53. Spiwak R, Lett R, Rwanyuma L, Logsetty S. Creation of a standardized burn course for Low Income Countries: Meeting local needs. *Burns*. 2014;40(7):1292-9.
54. Mbwele B, Ide NL, Mrema JG, Sarah APW, Melnick JA, Manongi R. Learning from Health Care Workers' Opinions for Improving Quality of neonatal Health Care in Kilimanjaro region, northeast Tanzania. *Ann Med Health Sci Res*. 2014;4(1):105-14.
55. Atreya MR, Lorenz JM, Narendran V. Provider Perceptions of Bubble Continuous Positive Airway Pressure and Barriers to Implementation in a Level III Neonatal Unit in South India. *Adv Neonatal Care*. 2018;18(6):500-6.

56. Bana S, Punjwani R, Khatoon A. Capacity building strategies for critical pediatric oncology nursing in a limited resource setting: A report of Children Cancer Hospital, Karachi, Pakistan. *Pediatric Blood and Cancer*. 2015;62(Supplement 4):S190.
57. Kandeel NA, Attia AK. Physical restraints practice in adult intensive care units in Egypt. *Nursing & Health Sciences*. 2013;15(1):79-85.
58. Munyiginya P, Brysiewicz P, Mill J. Critical care nursing practice and education in Rwanda. *South Afr J Crit Care*. 2016;32(2):55-7.
59. Saria VF, Mselle LT, Sicheloff BA. Parents and nurses telling their stories: the perceived needs of parents caring for critically ill children at the Kilimanjaro Christian Medical Centre in Tanzania. *BMC Nursing*. 2019;18(1).

Bibliography of citations in Table 3

1. Kassew T, Tilahun AD, Liyew B. Nurses' Knowledge, Attitude, and Influencing Factors regarding Physical Restraint Use in the Intensive Care Unit: A Multicenter Cross-Sectional Study. *Crit Care Res Pract*. 2020;2020.
2. Okech UK, Chokwe T, Mung'ayi V. The operational setup of intensive care units in a low income country in East Africa. *EAST AFRICAN MEDICAL JOURNAL*. 2015;92(2).
3. Gundo R. Contextual issues that influence preparedness of nurses for critical care nursing practice in Malawi. *Malawi Med J*. 2019;31(2):138-43.
4. Phuma-Ngaiyaye E, Kalembo FW. Supporting mothers to bond with their newborn babies: Strategies used in a neonatal intensive care unit at a tertiary hospital in Malawi. *International Journal of Nursing Sciences*. 2016;3(4):362-6.
5. Bruce JC, Dippenaar J, Schmollgruber S, Mphuthi DD, Huiskamp A. Advancing nursing scholarship: the Mozambique model. *Global Health Action*. 2017;10:1351116.
6. Ilesanmi RE, Olabisi P. Assessment of Common Interventions and Perceived Barriers to Pressure Ulcer Prevention in Southwest Nigeria. *J Wound Ostomy Cont Nurs*. 2014;41(3):242-6.
7. Shobowale EO, Onyedibe KI, Adegunle KB, Elikwu CJ. An Observational and Trend Analysis Study of Hand Hygiene Practices of Healthcare Workers at A Private Nigerian Tertiary Hospital. *Ann Med Health Sci Res*. 2017;7:84-9.
8. Ufashingabire CM, Nsereko E, Njunwa KJ, Brysiewicz P. Knowledge and attitudes of nurses regarding pain in the intensive care unit patients in Rwanda. *Rwanda Journal*. 2016;3(1):21-6.
9. Mbwele B, Ide NL, Mrema JG, Sarah APW, Melnick JA, Manongi R. Learning from Health Care Workers' Opinions for Improving Quality of neonatal Health Care in Kilimanjaro region, northeast Tanzania. *Ann Med Health Sci Res*. 2014;4(1):105-14.
10. Cunningham C, Brysiewicz P, Sepeku A, White L, Murray B, Lobue N, et al. Developing an emergency nursing short course in Tanzania. *Afr J Emerg Med*. 2017;7(4):147-50.
11. MacDonell R, Koe S, Okafor I, James A, Ahern C, Scanlan T. Building resuscitation capability in low resource settings; development of the PAIRS course (outlining the development of a tailored paediatric resuscitation programme for low resource healthcare settings). *Archives of Disease in Childhood*. 2019;104(Supplement 3):A259.
12. Spiwak R, Lett R, Rwanyuma L, Logsetty S. Creation of a standardized burn course for Low Income Countries: Meeting local needs. *Burns*. 2014;40(7):1292-9.
13. Byrne-Davis L, Slattery H, Whiteside S, Moro E, Jackson M, Onyachi N, et al. Efficacy and acceptability of an acute illness management course delivered to staff and students in Uganda by staff from the UK. *Special Issue: Disease elimination*. 2015;7(5):360-6.
14. Marzuola JM. Teaching Strategies to Reduce VAP at Mulago Hospital: A Capstone Project. 2016.
15. Kizza IB, Muliira JK. Nurses' pain assessment practices with critically ill adult patients. *International Nursing Review*. 2015;62(4):573-82.

16. Nakate GM, Emelonye AU. Examining nursing practices for management of sepsis in low income countries: the case of Uganda. *World*. 2018;12(1):2.
17. Carter C, Mukonka PS, Sitwala LJ, Howard-Hunt B, Notter J. The development of critical care nursing education in Zambia. *British journal of nursing* (Mark Allen Publishing). 2020;29(9):499-505.
18. Carter C, Snell D. Nursing the critically ill surgical patient in Zambia. *Br J Nurs*. 2016;25(20):1123-8.
19. Harrington J, Hall D, Wang TY, Murali M, Mwansa S. A quality improvement project to prevent hyperoxia in a developing world intensive care unit. *Critical Care*. 2020;24(Supplement 1).
20. . !!! INVALID CITATION !!! (20-29).
21. Pradhan N, Lama S, Mandal G, Shrestha E. Physical restraining: Nurses knowledge and practice in Tertiary Care Hospital of Eastern Nepal. *Nurs Open*. 2019;6(3):1029-37.
22. Niemczura J. Critical care nursing in nepal. *ANA Maine Journal*. 2015:8.
23. Hassan N, Afzal M, Sehar S, Gilani A. Effect of Educational Program on Pressure Ulcer Prevention Intervention among Nurses of Intensive Care Units at a Public Hospital, Pakistan. *Clinical Social Work and Health Intervention*. 2020;11(4):38-45.
24. Manasia RJM, Husain SJ, Hooda K, Imran M, Bailey C. Therapeutic Hypothermia Post-Cardiac Arrest A Clinical Nurse Specialist Initiative in Pakistan. *Clin Nurse Spec*. 2014;28(4):231-9.
25. Goonewardena CS, Colombage TD. Knowledge and practices of nurses caring for patients with endotracheal tube admitted to intensive care units in National Hospital of Sri Lanka. *Sri Lankan J Anaesthesiol*. 2020;28(2):94-100.
26. Koy V, Yunibhand J, Turale S. "It is really so exhausting": Exploring intensive care nurses' perceptions of 24-hour long shifts. *Journal of Clinical Nursing*.10.
27. Barkley AS, Medina-Beckwith J, Sothea S, Pak S, Durfy SJ, Lele AV. Teaching the Emergency Neurologic Life Support Course at Two Major Hospitals in Phnom Penh, Cambodia. *World neurosurgery*. 2020.
28. Neely J, Bashford T, Aldam P, Patel D, Ward L, Wilson J, et al. Developing sustainable Systems-Based Training in Trauma Intensive Care in Myanmar. *Journal of the Intensive Care Society*. 2018;19(2 Supplement 1):152-3.
29. Paguio JT, Banayat AC. Commentary on challenges to critical care nursing practice in the Philippines. *The World of Critical Care Nursing*. 2018;12(1).
30. Martinez RCKP, Rogado MIC, Serondo DJF, Soriano GP, Ilano KCS. Critical Care Nursing in the Philippines: Historical Past, Current Practices, and Future Directions. *Critical care nursing clinics of North America*. 2021;33(1):75-87.
31. Fadadu PP, Liu JC, Schiltz BM, Xoay TD, Phuc PH, Kumbamu A, et al. A Mixed-Methods Exploration of Pediatric Intensivists' Attitudes toward End-of-Life Care in Vietnam. *J Palliat Med*. 2019;22(8):885-93.
32. Emani SS, Allan CK, Forster T, Fisk AC, Lagrasta C, Zheleva B, et al. Simulation training improves team dynamics and performance in a low-resource cardiac intensive care unit. *Ann Pediatr Cardiol*. 2018;11(2):130-6.
33. Khallil NS. Critical care nurses' use of non-pharmacological pain management methods in Egypt. *Appl Nurs Res*. 2018;44:33-8.
34. Eltaybani S, Abdelwareth M, Abou-Zeid NA, Ahmed N. Recommendations to prevent nursing errors: Content analysis of semi-structured interviews with intensive care unit nurses in a developing country. *Journal of Nursing Management*. 2020;28(3):690-8.
35. AL-Gabri MM, Mohammed MA, Mehany MM. Factors Affecting Nurses Compliance Regarding the Care of Patients with Chest Trauma.

36. Al-Rafay SS, Al-Sharkawy SS. Educational Outcomes Associated With Providing a Comprehensive Guidelines Program About Nursing Care of Preterm Neonates Receiving Total Parenteral Nutrition. *Clin Nurs Res.* 2012;21(2):142-58.
37. Mohamed SS, Ali BAEM, Abd-El-Karim HE, Masoed ES. Nurses' knowledge and performance regarding infection control during milk formula preparation and its effect on neonates at El-Minia city hospitals. *The Journal of American Science.* 2012;8(10):782-7.
38. Zahran EM, Taha EE. Integrating evidence based nursing into the critical care nursing course: challenges from students' perspectives. *The Journal of American Science.* 2011;7(7):135-44.
39. Kandeel NA, Attia AK. Physical restraints practice in adult intensive care units in Egypt. *Nursing & Health Sciences.* 2013;15(1):79-85.
40. Zyoud SH, Khaled SM, Kawasmi BM, Habeba AM, Hamadneh AT, Anabosi HH, et al. Knowledge about the administration and regulation of high alert medications among nurses in Palestine: a cross-sectional study. *BMC Nursing.* 2019;18.
41. Guiles S, Lemons J, Trautman M, Bucher S, Songok J, Gisore P. The Implementation of a Neonatal Nurse Training Program at the Riley Mother Baby Hospital of Kenya. *Newborn Infant Nurs Rev.* 2016;16(4):184-9.
42. Kumar R, Canarie MF. Developing Pediatric Critical Care in Kenya*. *Pediatric Critical Care Medicine.* 2019;20(12):E538-E45.
43. Waweru-Siika W, Mung'ayi V, Misango D, Mogi A, Kisia A, Ngumi Z. The history of critical care in Kenya. *J Crit Care.* 2020;55:122-7.
44. Gundo R. Development, Implementation and Evaluation of an In-service Training Programme for Critical Care Nurses in Malawi. 2019.
45. Munyiginya P, Brysiewicz P, Mill J. Critical care nursing practice and education in Rwanda. *South Afr J Crit Care.* 2016;32(2):55-7.
46. Atumanya P, Sendagire C, Wabule A, Mukisa J, Ssemogerere L, Kwizera A, et al. Assessment of the current capacity of intensive care units in Uganda; A descriptive study. *Journal of Critical Care.* 2020;55:95-9.
47. Mwewa B, Mweemba P. Knowledge and utilization of ICU admission criteria and guidelines, Lusaka, Zambia. *Medical Journal of Zambia.* 2010;37(3):143-52.
48. Tadyanemhandu C, van Aswegen H, Ntsiea V. Early mobilisation practices of patients in intensive care units in Zimbabwean government hospitals - a cross-sectional study. *South Afr J Crit Care.* 2018;34(1):16-21.
49. Sahni N, Biswal M, Gandhi K, Kaur K, Saini V, Yaddanapudi LN. Effect of Intensive Education and Training of Nurses on Ventilator-associated Pneumonia and Central Line-associated Bloodstream Infection Incidence in Intensive Care Unit at a Tertiary Care Center in North India. *Indian Journal of Critical Care Medicine.* 2017;21(11):779-82.
50. Atreya MR, Lorenz JM, Narendran V. Provider Perceptions of Bubble Continuous Positive Airway Pressure and Barriers to Implementation in a Level III Neonatal Unit in South India. *Adv Neonatal Care.* 2018;18(6):500-6.
51. Rungta N, Zirpe KG, Dixit SB, Mehta Y, Chuadhry D, Govil D, et al. Indian Society of Critical Care Medicine Experts Committee Consensus Statement on ICU Planning and Designing, 2020. *Indian Journal of Critical Care Medicine.* 2020;24:S43-S60.
52. Khanal A, Sharma A, Basnet S. Current State of Pediatric Intensive Care and High Dependency Care in Nepal. *Pediatric Critical Care Medicine.* 2016;17(11):1032-40.
53. Limbu S, Kongsuwan W, Yodchai K. Lived experiences of intensive care nurses in caring for critically ill patients. *Nursing in Critical Care.* 2019;24(1):9-14.
54. Charlton S, O'Reilly G, Jones T, Fitzgerald M. Emergency care in developing nations: The role of emergency nurses in Galle, Sri Lanka. *Australas Emerg Nurs J.* 2011;14(2):69-74.

55. Haniffa R, De Silva AP, Iddagoda S, Batawalage H, De Silva S, Mahipala PG, et al. A cross-sectional survey of critical care services in Sri Lanka: A lower middle-income country. *Journal of Critical Care*. 2014;29(5):764-8.
56. Gautam P, Acharya SP, Williams G. Formation of the Southern Asia Association for Regional Cooperation Federation of Critical Care Nurses. *The World of Critical Care Nursing*. 2018;12(1).
57. Stephens T, De Silva AP, Beane A, Welch J, Sigera C, De Alwis S, et al. Capacity building for critical care training delivery: Development and evaluation of the Network for Improving Critical care Skills Training (NICST) programme in Sri Lanka. *Intensive & critical care nursing*. 2017;39:28-36.
58. Westcott M, Martiniuk ALC, Fowler RA, Adhikari NKJ, Dalipanda T. Critical care resources in the Solomon Islands: a cross-sectional survey. *BMC Int Health Hum Rights*. 2012;12:9.
59. Attia MAS, Youseff MRL, Abd El Fatah SAM, Ibrahim SK, Gomaa NAS. The relationship between health care providers' perceived work climate, organizational commitment, and caring efficacy at pediatric intensive care units, Cairo University. *Int J Health Plan Manag*. 2020;35(2):469-81.
60. Aloush SM, Abdelkader FA, Al-Sayaghi K, Tawalbeh LI, Suliman M, Al Bashtawy M, et al. Compliance of Nurses and Hospitals With Ventilator-Associated Pneumonia Prevention Guidelines: A Middle Eastern Survey. *Journal of Nursing Care Quality*. 2018;33(3):E8-E14.
61. MacLeod JBA, Jones T, Aphivantrakul P, Chupp M, Poenaru D. Evaluation of fundamental critical care course in Kenya: Knowledge, attitude, and practice. *Journal of Surgical Research*. 2011;167(2):223-30.
62. Sawe HR, Mfinanga JA, Lidenge SJ, Mpondo BCT, Msangi S, Lugazia E, et al. Disease patterns and clinical outcomes of patients admitted in intensive care units of tertiary referral hospitals of Tanzania. *BMC Int Health Hum Rights*. 2014;14.
63. Baker T, Lugazia E, Eriksen J, Mwafongo V, Irestedt L, Konrad D. Emergency and critical care services in Tanzania: a survey of ten hospitals. *BMC Health Services Research*. 2013;13.
64. Patel DV, Nimbalkar SM, Singh US, Kharod NM. Acceptability and implementation of FIMNCI by medical officers and staff nurses in government health institutions of Western India. *Archives of Disease in Childhood*. 2012;97(SUPPL. 2):A489-A90.
65. Sharma R, Jagota R, Koushal V. Study of Knowledge, Awareness and Practices of Infection Control among ICU Staff of a Multispecialty Tertiary Level Teaching Institute of North India. *J Health Manag*. 2013;15(1):45-56.
66. Paul R, McCutcheon SP, Zenios SA, Tregarthen JP, Denend LT. Sustaining Pressure Ulcer Best Practices in a High-Volume Cardiac Care Environment How one hospital reduced the incidence of hospital-acquired pressure ulcers to zero. *Am J Nurs*. 2014;114(8):34-44.
67. Swathi S, Ramesh A, Nagapoornima M, Fernandes LM, Jisina C, Rao PNS, et al. Sustaining a "culture of silence" in the neonatal intensive care unit during nonemergency situations: A grounded theory on ensuring adherence to behavioral modification to reduce noise levels. *Int J Qual Stud Health Well-Being*. 2014;9:1-10.
68. Rungta N, Zirpe KG, Dixit SB, Mehta Y, Chaudhry D, Govil D, et al. Indian Society of Critical Care Medicine Experts Committee Consensus Statement on ICU Planning and Designing, 2020. *Indian journal of critical care medicine : peer-reviewed, official publication of Indian Society of Critical Care Medicine*. 2020;24(Suppl 1):S43.

Bibliography of citations in Table 4

1. Gundo R. Contextual issues that influence preparedness of nurses for critical care nursing practice in Malawi. *Malawi Med J*. 2019;31(2):138-43.
2. Gundo R. Development, Implementation and Evaluation of an In-service Training Programme for Critical Care Nurses in Malawi. 2019.

3. Bruce JC, Dippenaar J, Schmollgruber S, Mphuthi DD, Huiskamp A. Advancing nursing scholarship: the Mozambique model. *Global Health Action*. 2017;10:1351116.
4. Nankundwa E, Brysiewicz P. Lived experiences of Rwandan ICU nurses caring for patients with a do-not-resuscitate order. *South Afr J Crit Care*. 2017;33(1):19-22.
5. Munyiginya P, Brysiewicz P, Mill J. Critical care nursing practice and education in Rwanda. *South Afr J Crit Care*. 2016;32(2):55-7.
6. Cunningham C, Brysiewicz P, Sepeku A, White L, Murray B, Lobue N, et al. Developing an emergency nursing short course in Tanzania. *Afr J Emerg Med*. 2017;7(4):147-50.
7. Kizza IB, Muliira JK. Nurses' pain assessment practices with critically ill adult patients. *International Nursing Review*. 2015;62(4):573-82.
8. Marzuola JM. Teaching Strategies to Reduce VAP at Mulago Hospital: A Capstone Project. 2016.
9. Carter C, Mukonka PS, Sitwala LJ, Howard-Hunt B, Notter J. The development of critical care nursing education in Zambia. *British journal of nursing* (Mark Allen Publishing). 2020;29(9):499-505.
10. Klopper H, Uys LR. The state of nursing and nursing education in Africa: a country-by-country review: Sigma Theta Tau; 2013.
11. Stephens T, De Silva AP, Beane A, Welch J, Sigera C, De Alwis S, et al. Capacity building for critical care training delivery: Development and evaluation of the Network for Improving Critical care Skills Training (NICST) programme in Sri Lanka. *Intensive & critical care nursing*. 2017;39:28-36.
12. Paguio JT, Banayat AC. Commentary on challenges to critical care nursing practice in the Philippines. *The World of Critical Care Nursing*. 2018;12(1).
13. Khallil NS. Critical care nurses' use of non-pharmacological pain management methods in Egypt. *Appl Nurs Res*. 2018;44:33-8.
14. Gorman LL, McDowell JRS. Identifying the needs of critical and acute cardiac care nurses within the first two years of practice in Egypt using a nominal group technique. *Nurse Education in Practice*. 2018;28:127-34.
15. Zyoud SH, Khaled SM, Kawasmi BM, Habeba AM, Hamadneh AT, Anabosi HH, et al. Knowledge about the administration and regulation of high alert medications among nurses in Palestine: a cross-sectional study. *Bmc Nursing*. 2019;18:17.

Bibliography of citations in Table 5

1. Kimani G, Maina F, Rithara S. Perception of palliative care course among other programs in a government learning institution. *Supportive Care in Cancer*. 2017;25(2 Supplement 1):S106.
2. Waweru-Siika W, Mung'ayi V, Misango D, Mogi A, Kisia A, Ngumi Z. The history of critical care in Kenya. *J Crit Care*. 2020;55:122-7.
3. Guiles S, Lemons J, Trautman M, Bucher S, Songok J, Gisore P. The Implementation of a Neonatal Nurse Training Program at the Riley Mother Baby Hospital of Kenya. *Newborn Infant Nurs Rev*. 2016;16(4):184-9.
4. Kumar R, Canarie MF. Developing Pediatric Critical Care in Kenya*. *Pediatric Critical Care Medicine*. 2019;20(12):E538-E45.
5. MacLeod JBA, Jones T, Aphivantrakul P, Chupp M, Poenaru D. Evaluation of fundamental critical care course in Kenya: Knowledge, attitude, and practice. *Journal of Surgical Research*. 2011;167(2):223-30.
6. Hernandez J, Kioko M, Adeyinka A, Pinto R, Mercredi G, Petrescu M, et al. Pediatric fundamentals of critical care support (PFCCS) course performance among staff from U.S. and resource-limited countries. *Pediatric Critical Care Medicine*. 2012;13(6):718.

7. Richards C, Fox J, Sheraton T. A questionnaire survey to aid development of a new critical care course in Liberia, West Africa. *European Journal of Anaesthesiology*. 2014;31(SUPPL. 52):242.
8. Gundo R. Development, Implementation and Evaluation of an In-service Training Programme for Critical Care Nurses in Malawi. 2019.
9. Bruce JC, Schmollgruber S, Baumann J. Inter-country master's degree in nursing: policy implications for the Mozambican health system. *International Nursing Review*. 2018;65(3):425-33.
10. Bruce JC, Dippenaar J, Schmollgruber S, Mphuthi DD, Huiskamp A. Advancing nursing scholarship: the Mozambique model. *Global Health Action*. 2017;10:1351116.
11. Munyiginya P, Brysiewicz P, Mill J. Critical care nursing practice and education in Rwanda. *South Afr J Crit Care*. 2016;32(2):55-7.
12. Mukamana D. Rwanda. In: Klopper H, Uys LR, editors. *The State of Nursing and Nursing Education in Africa: A Country-by-Country Review*. Indianapolis: Sigma Theta Tau International; 2013. p. 231-47.
13. Cunningham C, Brysiewicz P, Sepeku A, White L, Murray B, Lobue N, et al. Developing an emergency nursing short course in Tanzania. *Afr J Emerg Med*. 2017;7(4):147-50.
14. MacDonell R, Koe S, Okafor I, James A, Ahern C, Scanlan T. Building resuscitation capability in low resource settings; development of the PAIRS course (outlining the development of a tailored paediatric resuscitation programme for low resource healthcare settings). *Archives of Disease in Childhood*. 2019;104(Supplement 3):A259.
15. Mbwele B, Ide NL, Mrema JG, Sarah APW, Melnick JA, Manongi R. Learning from Health Care Workers' Opinions for Improving Quality of neonatal Health Care in Kilimanjaro region, northeast Tanzania. *Ann Med Health Sci Res*. 2014;4(1):105-14.
16. Carter C, Mukonka PS, Sitwala LJ, Howard-Hunt B, Notter J. The development of critical care nursing education in Zambia. *British journal of nursing (Mark Allen Publishing)*. 2020;29(9):499-505.
17. Klopper H, Uys LR. The state of nursing and nursing education in Africa: a country-by-country review: Sigma Theta Tau; 2013.
18. Patel DV, Nimbalkar SM, Singh US, Kharod NM. Acceptability and implementation of FIMNCI by medical officers and staff nurses in government health institutions of Western India. *Archives of Disease in Childhood*. 2012;97(SUPPL. 2):A489-A90.
19. Khanal A, Sharma A, Basnet S. Current State of Pediatric Intensive Care and High Dependency Care in Nepal. *Pediatric Critical Care Medicine*. 2016;17(11):1032-40.
20. Bana S, Punjwani R, Khatoon A. Capacity building strategies for critical pediatric oncology nursing in a limited resource setting: A report of Children Cancer Hospital, Karachi, Pakistan. *Pediatric Blood and Cancer*. 2015;62(Supplement 4):S190.
21. Stephens T, De Silva AP, Beane A, Welch J, Sigera C, De Alwis S, et al. Capacity building for critical care training delivery: Development and evaluation of the Network for Improving Critical care Skills Training (NICST) programme in Sri Lanka. *Intensive & critical care nursing*. 2017;39:28-36.
22. Haniffa R, De Silva AP, Iddagoda S, Batawalage H, De Silva S, Mahipala PG, et al. A cross-sectional survey of critical care services in Sri Lanka: A lower middle-income country. *Journal of Critical Care*. 2014;29(5):764-8.
23. Charlton S, O'Reilly G, Jones T, Fitzgerald M. Emergency care in developing nations: The role of emergency nurses in Galle, Sri Lanka. *Australas Emerg Nurs J*. 2011;14(2):69-74.
24. Gautam P, Acharya SP, Williams G. Formation of the Southern Asia Association for Regional Cooperation Federation of Critical Care Nurses. *The World of Critical Care Nursing*. 2018;12(1).

25. Neely J, Bashford T, Aldam P, Patel D, Ward L, Wilson J, et al. Developing sustainable Systems-Based Training in Trauma Intensive Care in Myanmar. *Journal of the Intensive Care Society*. 2018;19(2 Supplement 1):152-3.
26. Paguio JT, Banayat AC. Commentary on challenges to critical care nursing practice in the Philippines. *The World of Critical Care Nursing*. 2018;12(1).
27. Bechara AD, Tan M, Nguyen T, Le Doan A, Zheleva B. Creating professional development opportunities for pediatric cardiac critical care nurses in South Vietnam. *Cardiology in the Young*. 2017;27(4):S229.
28. Kandeel N, Ibrahim Y. Student Nurses' Perception on the Impact of Information Technology on Teaching and Learning. *Int J Inf Commun Technol Educ*. 2010;6(3):38-50.
29. Zahran EM, Taha EE. Integrating evidence based nursing into the critical care nursing course: challenges from students' perspectives. *The Journal of American Science*. 2011;7(7):135-44.
30. Zyoud SH, Khaled SM, Kawasmi BM, Habeba AM, Hamadneh AT, Anabosi HH, et al. Knowledge about the administration and regulation of high alert medications among nurses in Palestine: a cross-sectional study. *Bmc Nursing*. 2019;18.
31. Lutz J, Augustin M, Patel CC, Fenestor M, Patel A, Rock S, et al. IMPLEMENTATION OF A SUCCESSFUL INTENSIVE CARE NURSING STAFF DEVELOPMENT PROGRAM IN HAITI. *Critical Care Medicine*. 2014;42(12):1.
32. Gundo R. Development, Implementation and Evaluation of an In-Service Training Programme for Critical Care Nurses in Malawi 2019.
33. Ufashingabire CM, Nsereko E, Njunwa KJ, Brysiewicz P. Knowledge and attitudes of nurses regarding pain in the intensive care unit patients in Rwanda. *Rwanda Journal*. 2016;3(1):21-6.
34. Harrington J, Hall D, Wang TY, Murali M, Mwansa S. A quality improvement project to prevent hyperoxia in a developing world intensive care unit. *Critical Care*. 2020;24(Supplement 1).
35. Sathish Y, Lewis LE, Noronha JA, Nayak BS, Pai MS, Altimier L. Promoting developmental supportive care in preterm infants and families in a level III neonatal intensive care unit (NICU) setting in India. *Nurse Education in Practice*. 2019;40:8.
36. Atreya MR, Lorenz JM, Narendran V. Provider Perceptions of Bubble Continuous Positive Airway Pressure and Barriers to Implementation in a Level III Neonatal Unit in South India. *Adv Neonatal Care*. 2018;18(6):500-6.
37. Swathi S, Ramesh A, Nagapoornima M, Fernandes LM, Jisina C, Rao PNS, et al. Sustaining a "culture of silence" in the neonatal intensive care unit during nonemergency situations: A grounded theory on ensuring adherence to behavioral modification to reduce noise levels. *Int J Qual Stud Health Well-Being*. 2014;9:1-10.
38. Acharya R, Mishra SB, Ipsita S, Azim A. Impact of Nursing Education on CLABSI Rates: An Experience from a Tertiary Care Hospital in Eastern India. *Indian Journal of Critical Care Medicine*. 2019;23(7):316-9.
39. Dongara AR, Nimbalkar SM, Phatak AG, Patel DV, Nimbalkar AS. An Educational Intervention to Improve Nurses' Understanding of Pain in Children in Western India. *Pain Manag Nurs*. 2017;18(1):24-32.
40. Paul R, McCutcheon SP, Zenios SA, Tregarthen JP, Denend LT. Sustaining Pressure Ulcer Best Practices in a High-Volume Cardiac Care Environment How one hospital reduced the incidence of hospital-acquired pressure ulcers to zero. *Am J Nurs*. 2014;114(8):34-44.
41. Sahni N, Biswal M, Gandhi K, Kaur K, Saini V, Yaddanapudi LN. Effect of Intensive Education and Training of Nurses on Ventilator-associated Pneumonia and Central Line-associated Bloodstream Infection Incidence in Intensive Care Unit at a Tertiary Care Center in North India. *Indian Journal of Critical Care Medicine*. 2017;21(11):779-82.

42. Sharma R, Jagota R, Koushal V. Study of Knowledge, Awareness and Practices of Infection Control among ICU Staff of a Multispecialty Tertiary Level Teaching Institute of North India. *J Health Manag.* 2013;15(1):45-56.
43. Tripathy S, Routray PK, Mishra JC. Intensive Care Nurses' Attitude on Palliative and End of Life Care. *Indian Journal of Critical Care Medicine.* 2017;21(10):655-9.
44. Pradhan N, Lama S, Mandal G, Shrestha E. Physical restraining: Nurses knowledge and practice in Tertiary Care Hospital of Eastern Nepal. *Nurs Open.* 2019;6(3):1029-37.
45. Manasia RJM, Husain SJ, Hooda K, Imran M, Bailey C. Therapeutic Hypothermia Post-Cardiac Arrest A Clinical Nurse Specialist Initiative in Pakistan. *Clin Nurse Spec.* 2014;28(4):231-9.
46. Barkley AS, Medina-Beckwith J, Sothea S, Pak S, Durfy SJ, Lele AV. Teaching the Emergency Neurologic Life Support Course at Two Major Hospitals in Phnom Penh, Cambodia. *World neurosurgery.* 2020.
47. Emani SS, Allan CK, Forster T, Fisk AC, Lagrasta C, Zheleva B, et al. Simulation training improves team dynamics and performance in a low-resource cardiac intensive care unit. *Ann Pediatr Cardiol.* 2018;11(2):130-6.
48. Al-Gabri MM, Mohammed MA, Mehany MM. Factors Affecting Nurses Compliance Regarding the Care of Patients with Chest Trauma.
49. Al-Rafay SS, Al-Sharkawy SS. Educational Outcomes Associated With Providing a Comprehensive Guidelines Program About Nursing Care of Preterm Neonates Receiving Total Parenteral Nutrition. *Clin Nurs Res.* 2012;21(2):142-58.
50. Mohamed SS, Ali BAEM, Abd-El-Karim HE, Masoed ES. Nurses' knowledge and performance regarding infection control during milk formula preparation and its effect on neonates at El-Minia city hospitals. *The Journal of American Science.* 2012;8(10):782-7.
51. Eltaybani S, Abdelwareth M, Abou-Zeid NA, Ahmed N. Recommendations to prevent nursing errors: Content analysis of semi-structured interviews with intensive care unit nurses in a developing country. *Journal of Nursing Management.* 2020;28(3):690-8.
52. Aloush SM, Abdelkader FA, Al-Sayaghi K, Tawalbeh LI, Suliman M, Al Bashtawy M, et al. Compliance of Nurses and Hospitals With Ventilator-Associated Pneumonia Prevention Guidelines: A Middle Eastern Survey. *Journal of Nursing Care Quality.* 2018;33(3):E8-E14.
53. Ilesanmi RE, Olabisi P. Assessment of Common Interventions and Perceived Barriers to Pressure Ulcer Prevention in Southwest Nigeria. *J Wound Ostomy Cont Nurs.* 2014;41(3):242-6.
54. Byrne-Davis L, Slattery H, Whiteside S, Moro E, Jackson M, Onyachi N, et al. Efficacy and acceptability of an acute illness management course delivered to staff and students in Uganda by staff from the UK. *Special Issue: Disease elimination.* 2015;7(5):360-6.
55. Marzuola JM. Teaching Strategies to Reduce VAP at Mulago Hospital: A Capstone Project. 2016.
56. Bongar MVV, Pangan FC, Macindo JRB. Characteristics of a critical care clinical placement programme affecting critical care nursing competency of baccalaureate nursing students: A structural equation modelling. *Journal of Clinical Nursing.* 2019;28(9-10):1760-70.
57. Macindo JRB, Danganan MPB, Soriano CAF, Kho NSR, Bongar MVV. A conjoint analysis of the acute and critical care experiential learning preferences of Baccalaureate student nurses. *Nurse Education in Practice.* 2019;36:125-31.
58. Gorman LL, McDowell JRS. Identifying the needs of critical and acute cardiac care nurses within the first two years of practice in Egypt using a nominal group technique. *Nurse Education in Practice.* 2018;28:127-34.
59. Losonczy LI, Barnes SL, Liu S, Williams SR, McCurdy MT, Lemos V, et al. Critical care capacity in Haiti: A nationwide cross-sectional survey. *PLoS One.* 2019;14(6):e0218141.
60. Macleod JBA, Jones T, Aphivantrakul P, Chupp M, Poenaru D. Evaluation of fundamental critical care course in Kenya: knowledge, attitude, and practice. *The Journal of surgical research.* 2011;167(2):223-30.

61. Gundo R. Development, Implementation and Evaluation of an In-service Training Programme for Critical Care Nurses in Malawi: Auckland University of Technology; 2019.
62. Spiwak R, Lett R, Rwanyuma L, Logsetty S. Creation of a standardized burn course for Low Income Countries: Meeting local needs. Burns. 2014;40(7):1292-9.
63. Byrne-Davis L, Slattery H, Whiteside S, Moro E, Jackson M, Onyachi N, et al. Efficacy and acceptability of an acute illness management course delivered to staff and students in Uganda by staff from the UK. Int Health. 2015;7(5):360-6.

PRE PRINT